



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 


# PRODUCT INFORMATION



## (±)8(9)-EET

Item No. 50351

<b>Formal Name:</b>	(±)8,9-epoxy-5Z,11Z,14Z-eicosatrienoic acid
<b>Synonym:</b>	(±)8,9-EpETrE
<b>MF:</b>	C <sub>20</sub> H <sub>32</sub> O <sub>3</sub>
<b>FW:</b>	320.5
<b>Purity:</b>	≥98%
<b>Supplied as:</b>	A solution in ethanol
<b>Storage:</b>	-20°C
<b>Stability:</b>	≥2 years



NOTE: Relative stereochemistry shown in chemical structure

Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.

### Laboratory Procedures

(±)8(9)-EET is supplied as a solution in ethanol. To change the solvent, simply evaporate the ethanol under a gentle stream of nitrogen and immediately add the solvent of choice. Solvents such as DMSO and dimethyl formamide purged with an inert gas can be used. The solubility of (±)8(9)-EET in these solvents is approximately 50 mg/ml.

Further dilutions of the stock solution into aqueous buffers or isotonic saline should be made prior to performing biological experiments. Ensure that the residual amount of organic solvent is insignificant, since organic solvents may have physiological effects at low concentrations. If an organic solvent-free solution of (±)8(9)-EET is needed, it can be prepared by evaporating the ethanol and directly dissolving the neat oil in aqueous buffers. The solubility of (±)8(9)-EET in PBS (pH 7.2) is approximately 1 mg/ml. We do not recommend storing the aqueous solution for more than one day.

### Description

(±)8(9)-EET is a fully racemic version of the R/S enantiomeric forms biosynthesized from arachidonic acid (Item No. 90010).<sup>1,2</sup> (±)8(9)-EET is a major cytochrome P450 (CYP450) metabolite in the renal cortex.<sup>3</sup> It activates PPAR $\alpha$  in HEK293 cells when used at a concentration of 1  $\mu$ M but inhibits NF- $\kappa$ B activity induced by IL-1 $\beta$  in a PPAR $\alpha$ -dependent and -independent fashion.<sup>4</sup> It is also a substrate of COX-1 and COX-2, as measured by oxygen consumption and product formation assays when used at a concentration of 50  $\mu$ M.<sup>5</sup> (8S,9R)-EET (1  $\mu$ g/kg), but not (8R,9S)-EET, reduces the glomerular filtration rate (GFR) through COX-dependent pre-glomerular vasoconstriction in rats.<sup>6</sup>

### References

1. Chacos, N., Falck, J.R., Wixtrom, C., *et al.* Novel epoxides formed during the liver cytochrome P-450 oxidation of arachidonic acid. *Biochem. Biophys. Res. Commun.* **104(3)**, 916-922 (1982).
2. Oliw, E.H., Guengerich, F.P., and Oates, J.A. Oxygenation of arachidonic acid by hepatic monooxygenases. Isolation and metabolism of four epoxide intermediates. *J. Biol. Chem.* **257(7)**, 3771-3781 (1982).
3. Zhang, J.Y., Prakash, C., Yamashita, K., *et al.* Regiospecific and enantioselective metabolism of 8,9-epoxyeicosatrienoic acid by cyclooxygenase. *Biochem. Biophys. Res. Commun.* **183(1)**, 138-143 (1992).
4. Bystrom, J., Wray, J.A., Sugden, M.C., *et al.* Endogenous epoxygenases are modulators of monocyte/macrophage activity. *PLoS One* **6(10)**, e26591 (2011).
5. Rand, A.A., Barnych, B., Morisseau, C., *et al.* Cyclooxygenase-derived proangiogenic metabolites of epoxyeicosatrienoic acids. *Proc. Natl. Acad. Sci. USA* **114(17)**, 4370-4375 (2017).
6. Katoh, T., Takahashi, K., Capdevila, J., *et al.* Glomerular stereospecific synthesis and hemodynamic actions of 8,9-epoxyeicosatrienoic acid in rat kidney. *Am. J. Physiol.* **261(4 Pt 2)**, F578-F586 (1991).

#### WARNING

THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

#### SAFETY DATA

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

#### WARRANTY AND LIMITATION OF REMEDY

Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 05/09/2018

#### CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD  
ANN ARBOR, MI 48108 · USA

PHONE: [800] 364-9897  
[734] 971-3335

FAX: [734] 971-3640

CUSTSERV@CAYMANCHEM.COM  
WWW.CAYMANCHEM.COM