

# Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



## Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

## Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

## SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in



# PRODUCT INFORMATION



## Hexanolamino PAF C-16

Item No. 60905

CAS Registry No.: 137566-83-7

Formal Name: 1-O-hexadecyl-2-O-acetyl-sn-

glyceryl-3-phosphoryl(N,N,N-


trimethyl)hexanolamine

MF:  $C_{30}H_{62}NO_{7}P$ FW: 579.8 **Purity:** ≥98%

Supplied as: A lyophilized powder

Storage: -20°C Stability: ≥1 vear

Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.



### **Laboratory Procedures**

Hexanolamino PAF C-16 is supplied as a lyophilized powder. A stock solution may be made by dissolving the hexanolamino PAF C-16 in the solvent of choice, which should be purged with an inert gas. Hexanolamino PAF C-16 is soluble in organic solvents such as ethanol, DMSO, and dimethyl formamide. The solubility of hexanolamino PAF C-16 in these solvents is approximately 23, 0.5, and 1.5 mg/ml respectively.

Further dilutions of the stock solution into aqueous buffers or isotonic saline should be made prior to performing biological experiments. Ensure that the residual amount of organic solvent is insignificant, since organic solvents may have physiological effects at low concentrations. Organic solvent-free aqueous solutions of hexanolamino PAF C-16 can be prepared by directly dissolving the crystalline compound in aqueous buffers. The solubility of hexanolamino PAF C-16 in PBS (pH 7.2) is approximately 0.5 mg/ml. We do not recommend storing the aqueous solution for more than one day.

#### Description

Hexanolamino PAF C-16 is a PAF analog with mixed agonist/antagonist properties. In human monocyte-derived macrophages, it is an antagonist which inhibits the production of reactive oxygen species in response to PAF C-16.1 On the other hand, in rabbit platelets and guinea pig macrophages, hexanolamino PAF C-16 is a partial agonist, and in guinea pig platelets it is a full agonist.<sup>2</sup>

#### References

- 1. Rouis, M., Nigon, F., and Chapman, M.J. Platelet activating factor is a potent stimulant of the production of active oxygen species by human monocyte-derived macrophages. Biochem. Biophys. Res. Commun. 156(3), 1293-1301 (1988).
- 2. Stewart, A.G. and Grigoriadis, G. Structure-activity relationships for platelet-activating factor (PAF) and analogs reveal differences between PAF receptors on platelets and macrophages. J. Lipid Mediat. 4(3), 299-308 (1991).

WARNING
THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

## WARRANTY AND LIMITATION OF REMEDY

subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website

Copyright Cayman Chemical Company, 11/11/2021

### **CAYMAN CHEMICAL**

1180 EAST ELLSWORTH RD ANN ARBOR, MI 48108 · USA PHONE: [800] 364-9897

[734] 971-3335

FAX: [734] 971-3640 CUSTSERV@CAYMANCHEM.COM WWW.**CAYMANCHEM**.COM