



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 


# PRODUCT INFORMATION



## 1,2-Dioleoyl-*sn*-glycerol

Item No. 62230

CAS Registry No.: 24529-88-2  
Formal Name: 1,2-bis(O-9Z-octadecenoyl)-*sn*-glycerol  
MF: C<sub>39</sub>H<sub>72</sub>O<sub>5</sub>  
FW: 621.0  
Purity: ≥95%  
Supplied as: A solution in acetonitrile  
Storage: -20°C  
Stability: ≥1 year



Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.

### Laboratory Procedures

1,2-Dioleoyl-*sn*-glycerol is supplied as a solution in acetonitrile. To change the solvent, simply evaporate the acetonitrile under a gentle stream of nitrogen and immediately add the solvent of choice. Solvents such as ethanol, DMSO, and dimethyl formamide purged with an inert gas can be used. The solubility of 1,2-dioleoyl-*sn*-glycerol in these solvents is approximately 3, 7, and 20 mg/ml, respectively. 1,2-Dioleoyl-*sn*-glycerol is stable for at least six months in these solvents if stored at -20°C.

Further dilutions of the stock solution into aqueous buffers or isotonic saline should be made prior to performing biological experiments. Ensure that the residual amount of organic solvent is insignificant, since organic solvents may have physiological effects at low concentrations. If an organic solvent-free solution of 1,2-dioleoyl-*sn*-glycerol is needed, it can be prepared by evaporating the acetonitrile and directly dissolving the neat oil in aqueous buffers. The solubility of 1,2-dioleoyl-*sn*-glycerol in PBS (pH 7.2) is approximately 250 µg/ml. We do not recommend storing the aqueous solution for more than one day.

### Description

1,2-Dioleoyl-*sn*-glycerol is one of the PKC-activating second messengers collectively called diacyl glycerols (DAG). 1,2-Dioleoyl-*sn*-glycerol and 1,2-dioctanoyl-*sn*-glycerol were nearly equipotent in induction of the acrosome reaction in human sperm.<sup>1</sup>

### References

1. Doherty, C.M., Tarchala, S.M., Radwanska, E., *et al.* Characterization of two second messenger pathways and their interactions in eliciting the human sperm acrosome reaction. *J. Androl.* **16**, 36-46 (1995).

#### WARNING

THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

#### SAFETY DATA

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

#### WARRANTY AND LIMITATION OF REMEDY

Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 05/11/2018

#### CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD  
ANN ARBOR, MI 48108 · USA

PHONE: [800] 364-9897  
[734] 971-3335

FAX: [734] 971-3640

CUSTSERV@CAYMANCHEM.COM  
WWW.CAYMANCHEM.COM