



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic)


PRODUCT INFORMATION



C8 Ceramine (d18:1/8:0)

Item No. 62545

CAS Registry No.: 170926-06-4
Formal Name: N-octylsphingosine
Synonyms: C8 Ceramine, Ceramine (d18:1/8:0)
MF: C₂₆H₅₃NO₂
FW: 411.7
Purity: ≥98%
Supplied as: A crystalline solid
Storage: -20°C
Stability: ≥1 year



Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.

Laboratory Procedures

C8 Ceramine is supplied as a crystalline solid. A stock solution may be made by dissolving the C8 ceramine in the solvent of choice. C8 Ceramine is soluble in organic solvents such as ethanol, DMSO, and dimethyl formamide, which should be purged with an inert gas. The solubility of C8 ceramine in these solvents is approximately 33, 20, and 22 mg/ml, respectively.

C8 Ceramine is sparingly soluble in aqueous buffers. For maximum solubility in aqueous buffers, C8 ceramine should first be dissolved in ethanol and then diluted with the aqueous buffer of choice. C8 Ceramine has a solubility of approximately 0.5 mg/ml in a 1:1 solution of ethanol:PBS (pH 7.2) using this method. We do not recommend storing the aqueous solution for more than one day.

Description

C8 Ceramine is an analog of C8 ceramide (Item No. 62540) in which the carbonyl group of the ceramide is replaced by a methylene group. At a concentration of 10 μM, C8 ceramine induces maximal DNA fragmentation in U937 cells after 6 hours of incubation, compared to 12 hours for C8 ceramide.¹

Reference

1. Bernert, J.T., Jr., Bell, C.J., McGuffey, J.E., *et al.* Determination of "free" glycerol in human serum reference materials by isotope-dilution gas chromatography-mass spectrometry. *Journal of Chromatography* **578(1)**, 1-7 (1992).

WARNING
THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

SAFETY DATA
This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the [complete](#) Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY
Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 02/20/2018

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD
ANN ARBOR, MI 48108 · USA

PHONE: [800] 364-9897
[734] 971-3335

FAX: [734] 971-3640

CUSTSERV@CAYMANCHEM.COM
WWW.CAYMANCHEM.COM