



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

PRODUCT INFORMATION




T0901317

Item No. 71810

CAS Registry No.: 293754-55-9
Formal Name: N-(2,2,2-trifluoroethyl)-N-[4-[2,2,2-trifluoro-1-hydroxy-1-(trifluoromethyl)ethyl]phenyl]-benzenesulfonamide

MF: C₁₇H₁₂F₉NO₃S
FW: 481.3
Purity: ≥98%
UV/Vis.: λ_{max}: 227, 265, 272 nm
Supplied as: A crystalline solid
Storage: -20°C
Stability: ≥1 year



Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.

Laboratory Procedures

T0901317 is supplied as a crystalline solid. A stock solution may be made by dissolving the T0901317 in the solvent of choice, which should be purged with an inert gas. T0901317 is soluble in organic solvents such as ethanol, DMSO, and dimethyl formamide (DMF). The solubility of T0901317 in ethanol is approximately 50 mg/ml and approximately 100 mg/ml in DMSO and DMF.

T0901317 is sparingly soluble in aqueous buffers. For maximum solubility in aqueous buffers, T0901317 should first be dissolved in DMSO and then diluted with the aqueous buffer of choice. T0901317 has a solubility of approximately 0.5 mg/ml in a 1:1 solution of DMSO:PBS (pH 7.2) using this method. We do not recommend storing the aqueous solution for more than one day.

Description

T0901317 is a potent and selective agonist for both LXR α and LXR β , with an EC₅₀ of about 50 nM.¹ T0901317 acting through LXR and in concert with its RXR heterodimerization partner induces the expression of the ABCA1 reverse cholesterol transporter. This acts to increase the efflux of cholesterol from enterocytes and thus inhibit the overall absorption of cholesterol.

Reference

1. Repa, J.J., Turley, S.D., Lobaccaro, J.-M.A., *et al.* Regulation of absorption and ABC1-mediated efflux of cholesterol by RXR heterodimers. *Science* **289**, 1524-1529 (2000).

WARNING

THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

SAFETY DATA

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 01/08/2021

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD
ANN ARBOR, MI 48108 · USA

PHONE: [800] 364-9897
[734] 971-3335

FAX: [734] 971-3640

CUSTSERV@CAYMANCHEM.COM
WWW.CAYMANCHEM.COM