

Produktinformation



Forschungsprodukte & Biochemikalien
Zellkultur & Verbrauchsmaterial
Diagnostik & molekulare Diagnostik
Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien T. +43(0)1 489 3961-0 F. +43(0)1 489 3961-7 <u>mail@szabo-scandic.com</u> www.szabo-scandic.com

PRODUCT INFORMATION



(±)-Jasmonic Acid

Item No. 88300

CAS Registry No.: Formal Name:	77026-92-7 3-oxo-2R-(2Z)2-penten-1R-yl-cyclopentaneacetic acid	0
MF:	C ₁₂ H ₁₈ O ₃	$\wedge \sim - \sim$
FW:	210.3	$\langle \gamma \rangle \sim \langle \gamma \rangle$
Purity:	≥98% (sum of isomers)	Соон
Supplied as:	A solution in ethanol	
Storage:	-20°C	
Stability:	As supplied, 1 year from the QC date provided on the 0 stored properly	Certificate of Analysis, when

Laboratory Procedures

(±)-Jasmonic acid is supplied as a solution in ethanol. To change the solvent, simply evaporate the ethanol under a gentle stream of nitrogen and immediately add the solvent of choice. Solvents such as DMSO and dimethyl formamide (DMF) purged with an inert gas can be used. The solubility of (±)-jasmonic acid in DMF is approximately 25 mg/ml and approximately 16 mg/ml in DMSO.

Further dilutions of the stock solution into aqueous buffers or isotonic saline should be made prior to performing biological experiments. Ensure that the residual amount of organic solvent is insignificant, since organic solvents may have physiological effects at low concentrations. If an organic solvent-free solution of (±)-jasmonic acid is needed, it can be prepared by evaporating the ethanol and directly dissolving the neat oil in aqueous buffers. The solubility of (±)-jasmonic acid in PBS, pH 7.2, is approximately 3 mg/ml. We do not recommend storing the aqueous solution for more than one day.

Description

(±)-Jasmonic acid is a plant growth regulator and a derivative of α-linolenic acid (Item Nos. 90210 | 21910).¹ It decreases chlorophyll levels in green and etiolated barley leaf segments when used at concentrations of 10 and 50 μ M.² (±)-Jasmonic acid (10-1,000 μ M) inhibits elongation of rice seedlings.³ It increases germination of S. hermonthica seeds by 26% when used at a concentration of 1,000 μ M.⁴

References

- 1. Creelman, R.A. and Mullet, J.E. Biosynthesis and action of jasmonates in plants. Annu. Rev. Plant Physiol. Plant Mol. Biol. 48, 355-381 (1997).
- 2. Miersch, O., Herrmann, H., Kramell, H.-M., et al. Biological activity of jasmonic acid glucosyl ester. J. Plant Physiol. 182, 425-428 (1987).
- 3. Yamane, H., Sugawara, J., Suzuki, Y., et al. Synthesis of jasmonic acid related compounds and their structure-activity relationships on the growth of rice seedings. Agr. Biol. Chem. 44(12), 2857-2864 (1980).
- Yoneyama, K., Ogasawara, M., Takeuchi, Y., et al. Effect of jasmonates and related copmonuds on seed germination of Orobanche minor smith and Striga hermonthica (del.) benth. Biosci. Biotechnol. Biochem. 62(7), 1448-1450 (1998).

WARNING THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

SAFFTY DATA

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 01/07/2020

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD ANN ARBOR, MI 48108 · USA PHONE: [800] 364-9897 [734] 971-3335 FAX: [734] 971-3640 CUSTSERV@CAYMANCHEM.COM WWW.CAYMANCHEM.COM