



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

PRODUCT INFORMATION




Nitrotyrosine BSA

Item No. 89542

Overview and Properties

Contents:	This vial contains 200 µg of nitrotyrosine BSA
Synonyms:	3-Nitrotyrosine, NT-BSA, Peroxynitrite-treated BSA
Form:	Liquid
Storage:	-80°C (as supplied)
Stability:	≥3 years
Storage Buffer:	500 µl PBS, pH 7.2
Application:	Western blot (WB); the recommended starting concentration for use as a positive control is 50 ng. Other applications were not tested, therefore optimal working concentration/dilution should be determined empirically.

Image



Lane 1: BSA (1 µg)
Lane 2: Nitrotyrosine BSA (0.01 µg)
Lane 3: Nitrotyrosine BSA (0.025 µg)
Lane 4: Nitrotyrosine BSA (0.05 µg)
Lane 5: Nitrotyrosine BSA (0.1 µg)
Lane 6: Nitrotyrosine BSA (0.2 µg)

WARNING
THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

SAFETY DATA
This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY
Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 01/12/2022

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD

ANN ARBOR, MI 48108 · USA

PHONE: [800] 364-9897

[734] 971-3335

FAX: [734] 971-3640

CUSTSERV@CAYMANCHEM.COM

WWW.CAYMANCHEM.COM

PRODUCT INFORMATION



Description

Nitrotyrosine is a post-translational modification that is formed by the nitration of tyrosine.¹ Under conditions of oxidative stress, tyrosine is oxidized by reactive oxygen species (ROS) or, in the presence of hydrogen peroxide and nitrite, by myeloperoxidase (MPO) to yield a tyrosine radical that reacts with reactive nitrogen species (RNS), such as nitric oxide or peroxynitrite, to form nitrotyrosine. It exists in a free or protein-bound form and is commonly used as a marker of nitrosative or oxidative stress.² Nitrotyrosine residues have been found in a variety of proteins, including LDL, surfactant protein A, angiotensin II, and human and bovine serum albumin.³ Free nitrotyrosine production induced by peroxynitrite is inhibited by the polyphenols epicatechin gallate, gallic acid, catechin, or epicatechin in cell-free assays, as well as in aortic rings isolated from normotensive and spontaneously hypertensive rats administered the antioxidant N-acetyl-cysteine (NAC; Item No. 20261).^{4,5} Nitrotyrosine levels are increased in the affected tissues of numerous pathological conditions, including atherosclerosis, cancer, ulcerative colitis, Alzheimer's disease, and Parkinson's disease.³ Autoantibodies that recognize nitrotyrosinated proteins are increased in the synovium of patients with rheumatoid arthritis and are positively correlated with joint and tendon inflammation.¹ This product can be used as a positive control for immunoblotting experiments using Cayman's Nitrotyrosine Polyclonal Antibody (Item No. 10189540) and Nitrotyrosine Monoclonal Antibody (Item No. 189542).

References

1. Smallwood, M.J., Nissim, A., Knight, A.R., *et al.* Oxidative stress in autoimmune rheumatic diseases. *Free Radic. Biol. Med.* **125**, 3-14 (2018).
2. Teixeira, D., Fernandes, R., Prudêncio, C., *et al.* 3-Nitrotyrosine quantification methods: Current concepts and future challenges. *Biochimie* **125**, 1-11 (2016).
3. Oldreive, C. and Rice-Evans, C. The mechanisms for nitration and nitrotyrosine formation *in vitro* and *in vivo*: Impact of diet. *Free Radic. Res.* **35(3)**, 215-231 (2001).
4. Pannala, A.S., Rice-Evans, C.A., Halliwell, B., *et al.* Inhibition of peroxynitrite-mediated tyrosine nitration by catechin polyphenols. *Biochem. Biophys. Res. Commun.* **232(1)**, 164-168 (1997).
5. Cabassi, A., Dumont, E.C., Girouard, H., *et al.* Effects of chronic N-acetylcysteine treatment on the actions of peroxynitrite on aortic vascular reactivity in hypertensive rats. *J. Hypertens.* **19(7)**, 1233-1244 (2001).

CAYMAN CHEMICAL
1180 EAST ELLSWORTH RD
ANN ARBOR, MI 48108 · USA
PHONE: [800] 364-9897
[734] 971-3335
FAX: [734] 971-3640
CUSTSERV@CAYMANCHEM.COM
WWW.CAYMANCHEM.COM