

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in



PRODUCT INFORMATION



Arachidonic Acid

Item No. 90010

CAS Registry No.: 506-32-1

Formal Name: 5Z,8Z,11Z,14Z-eicosatetraenoic acid

Synonym:

MF: $C_{20}H_{32}O_{2}$ FW: 304.5 **Purity:** ≥98%

Supplied as: A solution in ethanol

Storage: -20°C Stability: ≥2 years

Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.

Laboratory Procedures

Arachidonic acid is supplied as a solution in ethanol. To change the solvent, simply evaporate the ethanol under a gentle stream of nitrogen and immediately add the solvent of choice. Solvents such as DMSO or dimethyl formamide can be used. To prevent oxidation of arachidonic acid, the solvent should be purged with an inert gas. The solubility of arachidonic acid in these solvents is at least 100 mg/ml.

Further dilutions of the stock solution into aqueous buffers or isotonic saline should be made prior to performing biological experiments. Ensure that the residual amount of organic solvent is insignificant, since organic solvents may have physiological effects at low concentrations. If an organic solvent-free aqueous solution of arachidonic acid is needed, it can be prepared by evaporating the ethanol and directly dissolving the neat oil in basic buffers. Arachidonic acid is sparingly soluble in neutral buffers. Store aqueous solutions of arachidonic acid on ice and use within 12 hours of preparation. Although the aqueous solutions of arachidonic acid may be stable for more than 12 hours, we strongly recommend using a fresh preparation each day.

Description

Arachidonic acid is an essential fatty acid and a precursor for all prostaglandins, thromboxanes, and leukotrienes. Virtually all cellular arachidonic acid is esterified in membrane phospholipids where its presence is tightly regulated through multiple interconnected pathways.1 Free arachidonic acid is a transient, critical substrate for the biosynthesis of eicosanoid second messengers. Receptor-stimulated release, metabolism, and re-uptake of free arachidonate are all important aspects of cell signaling and inflammation.²

References

- 1. Nixon, A.B., Greene, D.G., and Wykle, R.L. Comparison of acceptor and donor substrates in the CoAindependent transacylase reaction in human neutrophils. Biochim. Biophys. Acta 1300, 187-196 (1996).
- 2. Burgoyne, R.D. and Morgan, A. The control of free arachidonic acid levels. Trends Biochem. Sci. 15, 365-366 (1990).

WARNING
THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

uyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 08/03/2017

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD ANN ARBOR, MI 48108 · USA PHONE: [800] 364-9897

[734] 971-3335

FAX: [734] 971-3640 CUSTSERV@CAYMANCHEM.COM WWW.**CAYMANCHEM**.COM