

Produktinformation



Forschungsprodukte & Biochemikalien
Zellkultur & Verbrauchsmaterial
Diagnostik & molekulare Diagnostik
Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien T. +43(0)1 489 3961-0 F. +43(0)1 489 3961-7 <u>mail@szabo-scandic.com</u> www.szabo-scandic.com

PRODUCT INFORMATION



Resolvin D2 methyl ester

Item No. 9001497

CAS Registry No.: Formal Name:	810668-53-2 7S,16R,17S-trihydroxy-		
Tormar Name.	4Z.8E.10Z.12E.14E.19Z-	HO.	
	docosahexaenoic acid, methyl ester		
Synonyms:	7(S),16(R),17(S)-Resolvin D2 methyl ester,		соосн ₃
	RvD2 methyl ester		
MF:	C ₂₃ H ₃₄ O ₅	ОН	
FW:	390.5		
Purity:	≥95%	\geq	
UV/Vis.:	λ _{max} : 302 nm	HO	
Supplied as:	A solution in ethanol		
Storage:	-80°C		
Stability:	≥1 year		
Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.			

Laboratory Procedures

Resolvin D2 (RvD2) methyl ester is supplied as a solution in ethanol. To change the solvent, simply evaporate the ethanol under a gentle stream of nitrogen and immediately add the solvent of choice. It is recommended that this product be stored and handled in an ethanol solution. Resolvins can isomerize and degrade when put into freeze thaw conditions and/or in solvents such as dimethyl formamide or DMSO.

Further dilutions of the stock solution into aqueous buffers or isotonic saline should be made prior to performing biological experiments. Ensure that the residual amount of organic solvent is insignificant, since organic solvents may have physiological effects at low concentrations. If an organic solvent-free solution of RvD2 methyl ester is needed, it can be prepared by evaporating the ethanol and directly dissolving the neat oil in aqueous buffers. The solubility of RvD2 methyl ester in PBS, pH 7.2, is approximately 0.05 mg/ml. Aqueous solutions of RvD2 methyl ester should be discarded immediately after use.

Description

RvD2 (Item No. 10007279) is a lipid mediator biosynthesized by the sequential oxygenation of docosahexaenoic acid by 15- and 5-lipoxygenase.¹ It evokes diverse anti-inflammatory effects which may mediate the resolution of inflammation.^{2,3} RvD2 methyl ester is a methyl ester version of the free acid which may act as a lipophilic prodrug form that will alter its distribution and pharmacokinetic properties. The methyl ester moiety is susceptible to cleavage by intracellular esterases, leaving the free acid.

References

- 1. Seki, H., Sasaki, T., Ueda, T., et al. Resolvins as regulators of the immune system. ScientificWorldJournal 10, 818-831 (2010).
- 2. Serhan, C.N. Novel lipid mediators and resolution mechanisms in acute inflammation: To resolve or not? Am. J. Pathol. 177(4), 1576-1591 (2011).
- 3. Spite, M., Norling, L.V., Summers, L., et al. Resolvin D2 is a potent regulator of leukocytes and controls microbial sepsis. Nature 461(7268), 1287-1291 (2009).

WARNING THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

SAFFTY DATA

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

uyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 06/14/2019

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD ANN ARBOR, MI 48108 · USA PHONE: [800] 364-9897 [734] 971-3335 FAX: [734] 971-3640 CUSTSERV@CAYMANCHEM.COM WWW.CAYMANCHEM.COM