

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in



PRODUCT INFORMATION



GSK805

Item No. 9002444

CAS Registry No.: 1426802-50-7

Formal Name: N-[2,6-dichloro-2'-(trifluoromethoxy)[1,1'-biphenyl]-4-

yl]-4-(ethylsulfonyl)-benzeneacetamide


MF: $C_{23}H_{18}CI_2F_3NO_4S$

FW: 532.4 **Purity:** ≥98%

 λ_{max} : 221, 256 nm A crystalline solid UV/Vis.: Supplied as:

-20°C Storage: Stability: ≥2 years

Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.



Laboratory Procedures

GSK805 is supplied as a crystalline solid. A stock solution may be made by dissolving the GSK805 in the solvent of choice. GSK805 is soluble in organic solvents such as DMSO and dimethyl formamide, which should be purged with an inert gas. The solubility of GSK805 in these solvents is approximately 5 and 3 mg/ml, respectively.

GSK805 is sparingly soluble in aqueous buffers. For maximum solubility in aqueous buffers, GSK805 should first be dissolved in DMSO and then diluted with the aqueous buffer of choice. GSK805 has a solubility of approximately 0.3 mg/ml in a 1:2 solution of DMSO:PBS (pH 7.2) using this method. We do not recommend storing the aqueous solution for more than one day.

Description

GSK805 is a potent, orally bioavailable retinoid-related orphan receptor gamma t (RORyt) inverse agonist that interacts with the receptor's putative ligand binding domain without exerting significant effects on DNA binding.¹ It inhibits the expression of IL-17 (at 0.5 μM) in naïve CD4⁺ T cells activated under Th17-cell-polarizing conditions and affects the broader RORyt-dependent gene network, inhibiting the development and pathogenic function of Th17 cells. GSK805 significantly reduces the severity of experimental autoimmune encephalomyelitis (EAE), a mouse model of multiple sclerosis, when given orally to the hosts at 10 mg/kg daily beginning at the time of disease induction.

Reference

1. Xiao, S., Yosef, N., Yang, J., et al. Small-molecule RORγt antagonists inhibit T helper 17 cell transcriptional network by divergent mechanisms. Immunity 40(4), 477-489 (2014).

WARNING
THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information Buyer agrees to purchase the material can be found on our website.

Copyright Cayman Chemical Company, 06/15/2017

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD ANN ARBOR, MI 48108 · USA PHONE: [800] 364-9897

[734] 971-3335

FAX: [734] 971-3640 CUSTSERV@CAYMANCHEM.COM WWW.**CAYMANCHEM**.COM