

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in



PRODUCT INFORMATION



Myristoleic Acid

Item No. 9002461

CAS Registry No.: 544-64-9

Formal Name: 9Z-tetradecenoic acid

Synonyms: cis-9-Tetradecenoate, Oleomyristic

MF: $C_{14}H_{26}O_{2}$ FW: 226.4 **Purity:** ≥95%

Supplied as: A solution in ethanol

Storage: -20°C Stability: ≥1 year

Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.

Laboratory Procedures

Myristoleic acid is supplied as a solution in ethanol. To change the solvent, simply evaporate the ethanol under a gentle stream of nitrogen and immediately add the solvent of choice. Solvents such as DMSO and dimethyl formamide purged with an inert gas can be used. The solubility of myristoleic acid in these solvents is approximately 2.5 mg/ml and 3 mg/ml, respectively.

Further dilutions of the stock solution into aqueous buffers or isotonic saline should be made prior to performing biological experiments. Ensure that the residual amount of organic solvent is insignificant, since organic solvents may have physiological effects at low concentrations. If an organic solvent-free solution of myristoleic acid is needed, it can be prepared by evaporating the ethanol and directly dissolving the neat oil in aqueous buffers. The solubility of myristoleic acid in PBS, pH 7.2, is approximately 2 mg/ml. We do not recommend storing the aqueous solution for more than one day.

Description

Myristoleic acid is a monounsaturated fatty acid found in the fruit extract of S. repens and in dairy extracts.^{1,2} It induces apoptosis and necrosis in human prostate cancer LNCaP cells at a rate of 8.8% and 8.1%, respectively. Myristoleic acid inhibits C. albicans germination in vitro with a minimal inhibitory concentration (MIC) of 9 µM.2 Furthermore, myristoleic acid inhibits osteogenesis in vitro via interference with cytoskeletal rearrangement and prevents RANKL-induced bone loss and osteoclast formation in mice.³

References

- 1. Iguchi, K., Okumura, N., Usui, S., et al. Myristoleic acid, a cytotoxic component in the extract from Serenoa repens, induces apoptosis and necrosis in human prostatic LNCaP cells. Prostate 47(1), 59-65 (2001).
- 2. Clément, M., Tremblay, J., Lange, M., et al. Whey-derived free fatty acids suppress the germination of Candida albicans in vitro. FEMS Yeast Res. 7(2), 276-285 (2007).
- Kwon, J.O., Jin, W.J., Kim, B., et al. Myristoleic acid inhibits osteoclast formation and bone resorption by suppressing the RANKL activation of Src and Pyk2. Eur. J. Pharmacol. 768, 189-198 (2015).

WARNING
THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information Buyer agrees to purchase the material can be found on our website.

Copyright Cayman Chemical Company, 02/14/2019

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD ANN ARBOR, MI 48108 · USA PHONE: [800] 364-9897

[734] 971-3335

FAX: [734] 971-3640 CUSTSERV@CAYMANCHEM.COM WWW.**CAYMANCHEM**.COM