



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com


[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

PRODUCT INFORMATION



Lignoceric Acid-d₄₇ Item No. 9003320

CAS Registry No.: 68060-00-4
Formal Name: tetracosanoic-d₄₇ acid
Synonym: C₂₄:0-d₄₇
MF: C₂₄H₄₇O₂
FW: 415.9
Chemical Purity: ≥98% (Lignoceric Acid)
Deuterium Incorporation: ≥99% deuterated forms (d₁-d₄₇); ≤1% d₀
Supplied as: A solid
Storage: -20°C
Stability: ≥2 years



Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.

Laboratory Procedures

Lignoceric acid-d₄₇ is intended for use as an internal standard for the quantification of lignoceric acid (Item No. 13353) by GC- or LC-MS. The accuracy of the sample weight in this vial is between 5% over and 2% under the amount shown on the vial. If better precision is required, the deuterated standard should be quantitated against a more precisely weighed unlabeled standard by constructing a standard curve of peak intensity ratios (deuterated versus unlabeled).

Lignoceric acid-d₄₇ is supplied as a solid. A stock solution may be made by dissolving the lignoceric acid-d₄₇ in the solvent of choice, which should be purged with an inert gas. Lignoceric acid-d₄₇ is soluble in organic solvents such as chloroform and THF. The solubility of lignoceric acid-d₄₇ in these solvents is approximately 2 and 5 mg/ml, respectively.

Description

Lignoceric acid is a 24-carbon saturated (24:0) fatty acid. In mammals, it is synthesized during brain development and is found in cerebroside.¹ The deficient peroxisomal oxidation of very-long-chain fatty acids, including lignoceric acid, contributes to certain syndromes, including Zellweger cerebro-hepato-renal syndrome and X chromosome-linked adrenoleukodystrophy.² Lignoceric acid is also a by-product of lignin production.

References

1. Bourre, J.M., Paturneau-Jouas, M.Y., Daudu, O.L., *et al.* Lignoceric acid biosynthesis in the developing brain. Activities of mitochondrial acetyl-CoA-dependent synthesis and microsomal malonyl-CoA chain-elongating system in relation to myelination. *Eur. J. Biochem.* **72(1)**, 41-47 (1977).
2. Singh, I., Moser, A.E., Goldfischer, S., *et al.* Lignoceric acid is oxidized in the peroxisome: Implications for the Zellweger cerebro-hepato-renal syndrome and adrenoleukodystrophy. *Proc. Natl. Acad. Sci. USA* **81(13)**, 4203-4207 (1984).

WARNING

THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

SAFETY DATA

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 12/17/2019

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD
ANN ARBOR, MI 48108 · USA

PHONE: [800] 364-9897
[734] 971-3335

FAX: [734] 971-3640

CUSTSERV@CAYMANCHEM.COM
WWW.CAYMANCHEM.COM