

Produktinformation



Forschungsprodukte & Biochemikalien
Zellkultur & Verbrauchsmaterial
Diagnostik & molekulare Diagnostik
Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien T. +43(0)1 489 3961-0 F. +43(0)1 489 3961-7 <u>mail@szabo-scandic.com</u> www.szabo-scandic.com

PRODUCT INFORMATION



Docosatrienoic Acid

Item No. 90170

CAS Registry No.:	28845-86-5	\frown \land \land \land \land
Formal Name:	13Z,16Z,19Z-docosatrienoic acid	/ / / Соон
Synonym:	cis-13,16,19-Docosatrienoic Acid	\langle
MF:	C ₂₂ H ₃₈ O ₂	
FW:	334.5	
Purity:	≥98%	
Supplied as:	A solution in ethanol	
Storage:	-20°C	
Stability:	≥1 year	
Special Conditions: Oxygen and light sensitive		
Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.		



Laboratory Procedures

Docosatrienoic acid is supplied as a solution in ethanol. To change the solvent, simply evaporate the ethanol under a gentle stream of nitrogen and immediately add the solvent of choice. Solvents such as DMSO and dimethyl formamide purged with an inert gas can be used. The solubility of docosatrienoic acid in these solvents is approximately 100 mg/ml. Docosatrienoic acid is also soluble in 0.15 M Tris-HCl, pH 8.5, at approximately 1 mg/ml.

Further dilutions of the stock solution into aqueous buffers or isotonic saline should be made prior to performing biological experiments. Ensure that the residual amount of organic solvent is insignificant, since organic solvents may have physiological effects at low concentrations. If an organic solvent-free solution of docosatrienoic acid is needed, it can be prepared by evaporating the ethanol and directly dissolving the neat oil in aqueous buffers. The solubility of docosatrienoic acid in PBS, pH 7.2, is approximately 100 μ g/ml. We do not recommend storing the aqueous solution for more than one day.

Description

Docosatrienoic acid is a rare ω -3 fatty acid not readily detected in normal phospholipid PUFA pools. It is a polyunsaturated version of the fatty acid docosanoic acid (Item No. 9000338). Docosatrienoic acid inhibits $[^{3}H]$ -LTB₄ binding to pig neutrophils at a concentration of 5 μ M.¹ It also dose-dependently inhibits the activity of mammalian DNA polymerases and human topoisomerases I and II, potentially by binding directly to the hydrophobic region of the enzymes.² The cis configuration of the double bonds is important for this activity, and the carboxyl group is required as the methyl ester version has no activity at these enzymes.

References

- 1. Yagaloff, K.A., Franco, L., Simko, B., et al. Essential fatty acids are antagonists of the leukotriene B4 receptor. Prostaglandins, Leukot. Essent. Fatty Acids 52(5), 293-297 (1995).
- 2. Yonezawa, Y., Hada, T., Uryu, K., et al. Inhibitory action of C22-fatty acids on DNA polymerases and DNA topoisomerases. Int. J. Mol. Med. 18(4), 583-588 (2006).

WARNING THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

SAFFTY DATA

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

uyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 09/18/2017

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD ANN ARBOR, MI 48108 · USA PHONE: [800] 364-9897 [734] 971-3335 FAX: [734] 971-3640 CUSTSERV@CAYMANCHEM.COM WWW.CAYMANCHEM.COM