



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

PRODUCT INFORMATION



α -Linolenoyl Ethanolamide

Item No. 90215

CAS Registry No.: 57086-93-8

Formal Name: N-(2-hydroxyethyl)-9Z,12Z,15Z-octadecatrienamide

Synonym: N-Linolenoyl Ethanolamine

MF: $C_{20}H_{35}NO_2$


FW: 321.5

Purity: $\geq 98\%$

Supplied as: A solution in ethanol

Storage: $-20^{\circ}C$

Stability: As supplied, 1 year from the QC date provided on the Certificate of Analysis, when stored properly



Laboratory Procedures

α -Linolenoyl ethanolamide is supplied as a solution in ethanol. To change the solvent, simply evaporate the ethanol under a gentle stream of nitrogen and immediately add the solvent of choice. Solvents such as DMSO and dimethyl formamide purged with an inert gas can be used. The solubility of α -linolenoyl ethanolamide in these solvents is approximately 30 and 10 mg/ml, respectively.

α -Linolenoyl ethanolamide is sparingly soluble in aqueous buffers. For maximum solubility in aqueous buffers, the ethanolic solution of α -linolenoyl ethanolamide should be diluted with the aqueous buffer of choice. α -Linolenoyl ethanolamide has a solubility of approximately 8.5 mg/ml in a 1:2 solution of ethanol:PBS (pH 7.2) using this method. We do not recommend storing the aqueous solution for more than one day.

Description

α -Linolenoyl ethanolamide is an endocannabinoid containing α -linolenic acid in place of the arachidonate moiety of AEA. α -Linolenoyl ethanolamide has been detected in porcine brain, but its specific role and relative importance as a cannabinergic neurotransmitter have not been elucidated.¹

Reference

1. Hanus, L., Gopher, A., Almog, S., *et al.* Two new unsaturated fatty acid ethanolamides in brain that bind to the cannabinoid receptor. *J. Med. Chem.* **36**, 3032-3034 (1993).

WARNING

THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

SAFETY DATA

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 06/28/2016

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD
ANN ARBOR, MI 48108 · USA

PHONE: [800] 364-9897

[734] 971-3335

FAX: [734] 971-3640

CUSTSERV@CAYMANCHEM.COM

WWW.CAYMANCHEM.COM