



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Technically  
Speaking

**CEDARLANE**<sup>®</sup>  
[www.cedarlanelabs.com](http://www.cedarlanelabs.com)



Conveniently Delivering You Today's Innovations  
for the Science of Tomorrow™

**Anti-Rat CD8a  
Monoclonal Antibody**

Catalogue#	Format	Size	Concentration	Isotype Control
<b>CL004A</b>	Ascites	0.5ml	N/A	CLCMG2A00
<b>CL004AP/-2</b>	Purified	250µg/500µg	1.0 mg/ml	CLCMG2A00
<b>CL004AP-S</b>	Purified	50µg	1.0 mg/ml	CLCMG2A00
<b>CL004B/-5</b>	Biotin	100µg/500µg	0.1 mg/ml	CLCMG2A15
<b>CL004F/-5</b>	FITC	100µg/500µg	0.1 mg/ml	CLCMG2A01
<b>CL004NA</b>	No Azide	1mg	1.0 mg/ml	CLCMG2A00
<b>CL004PE/-4</b>	PE	50µg/200µg	0.1 mg/ml	CLCMG2A04
<b>CL004TC</b>	PE-Cy5	100µg	0.1 mg/ml	CLCMG106
<b>CL004AF4</b>	Alexa Fluor <sup>®</sup> 488	100µg	0.1 mg/ml	N/A

Alexa Fluor<sup>®</sup> is a registered trademark of Life Technologies Corporation.

Isotype Control: Mouse IgG<sub>1</sub>

**DESCRIPTION:**

Cedarlane's anti-rat T cytotoxic/suppressor cell monoclonal antibody recognizes a determinant on the majority of thymocytes (90-95%), a subset of peripheral T cells, the majority of NK cells, and the granular intraepithelial leukocytes in the small intestine (1,2,3,4). The antigen recognized is a complex of surface glycoproteins of Mr 34, 39, and 76 kDa and is the rat homologue of the human CD8 and the mouse Ly 2.3 antigen (5,8). CL004 labels all peripheral T cells that are unlabeled by the CL003 (W3/25) monoclonal antibody. It labels a T cell subset which mediates the suppression of antibody formation (1) and the cytotoxic cell precursor (2). CL004 and CL003 can be used together to fractionate T cells by sorting in FACS or by rosette depletion (1,2) or can be used together to study subsets of T cells in the rat which mediate lethal graft versus host disease (7). This antibody is one of the 3 monoclonal antibodies which labels T lymphocyte populations in the rat, these being CL002AP (W3/13) which labels all T cells, as well as CL003 (W3/25) and CL004 (MRC OX-8) which label non-overlapping T cell populations. These monoclonal antibodies used in concert are being employed extensively to investigate cellular aspects of the immune response in rats and prove to be useful as markers for functionally distinct subpopulations of lymphocytes. This clone is reported to work with frozen and paraffin sections (9).

**PRESENTATION:**

**Ascites:** 0.5ml Lyophilized.

**Purified:** Purified IgG buffered in PBS and 0.02% NaN<sub>3</sub>. (Purified from ascitic fluid via Protein G Chromatography). For maximum recovery of contents, spin down tube before use.

**Biotin, FITC, PE and AF488:** Biotin/FITC/PE/AF488 conjugated IgG buffered in PBS, 0.02% NaN<sub>3</sub> and EIA grade BSA as a stabilizing protein to bring total protein concentration to 4-5 mg/ml.

Visit our website for your local distributor.

**CEDARLANE**<sup>®</sup>



[www.cedarlanelabs.com](http://www.cedarlanelabs.com)

An ISO 9001:2000 and ISO 13485:2003  
registered company.

In CANADA: Toll Free: 1-800-268-5058

4410 Paletta Court, Burlington, ON L7L 5R2 ph: (289) 288-0001, fax: (289) 288-0020  
e-mail: [general@cedarlanelabs.com](mailto:general@cedarlanelabs.com)

In the USA: Toll Free: 1-800-721-1644

1210 Turrentine Street, Burlington, NC 27215 ph: (336) 513-5135, fax: (336) 513-5138  
e-mail: [service@cedarlanelabs.com](mailto:service@cedarlanelabs.com)

## **STORAGE/STABILITY:**

Stable at 4°C. DO NOT FREEZE PE and AF488 conjugates For long term storage, aliquot and freeze unused portions at -20°C in volumes appropriate for single usage. Avoid repeated freeze thaw cycles.

## **SPECIFICATIONS:**

Clone: MRC OX-8

Hybridoma Production:

Immunization:

Immunogen: High molecular weight rat thymocyte glycoproteins

Donor: BALB/c spleen

Fusion Partner: P3-NSI-1-Ag4 (NS1/1)

Specificity: Rat CD8a

Strain Distributions:

Strains Tested: Wistar, Buffalo, Brown Norway, Fischer 344

Positive: Wistar, Buffalo, Brown Norway, Fischer 344

Negative: none

## **FLOW CYTOMETRIC ANALYSIS:**

Rat Strain: Wistar

Cell Concentration:  $1 \times 10^6$  cells per test

Antibody Concentration:  $0.1 \mu\text{g}/10^6$  cells

### **Cell source**

Thymus

Spleen


Lymph Node

### **Percent staining**

92.6%

11.8%

23.6%



Wistar rat splenic T-cells (left) or thymocytes (right) were stained with anti-CD8a (clone: OX-8) (filled histogram) or mouse IgG1,  $\kappa$  isotype control (open histogram).

**N.B.** Appropriate control samples should always be included in any labelling studies.

\* For optimal results in various applications, it is recommended that each investigator determine dilutions appropriate for individual use.

## **REFERENCES:**

1. Brideau, R.J., Carter, P.B., McMaster, W.R., Mason, D.W. and A.F. Williams. (1980) Eur. J. of Immunol. 10, 609-615. Two subsets of rat T lymphocytes defined with monoclonal antibodies.
2. Dallman, M.J., Mason, D.W. and M. Webb. (1982) Eur. J. Immunol. 12, 511-518. The role of host and donor cells in the rejection of skin allografts by T cell deprived rats infected with syngenic T cells.
3. Cantrell, D.A., Robins, R.A., Brooks, C.G. and R.W. Baldwin. (1982) Immunology. 45, 97-103. Phenotype of rat natural killer cells defined by monoclonal antibodies marking rat lymphocyte subsets.
4. Lyscom, N. and M.J. Brueton. (1982) Immunology. 45, 775-783. Intraepithelial lamina propria and Peyer's patch lymphocytes of the rat small intestine: isolation and characterization in terms of immunoglobulin markers and receptors for monoclonal antibodies.
5. Johnson, P., Gagnon, J., Barclay, A.N. and A.F. Williams. (1985) EMBO J. 4, 2539-2545. Purification, chain separation and sequence of the MRC OX-8 antigen, a marker of rat cytotoxic T lymphocytes.
6. Barclay, A.N. (1981) Immunology, 42, 593-600. The localization of populations of lymphocytes defined by monoclonal antibodies in rat lymphoid tissues.
7. Mason, D.W. (1981) Transplantation, 32, 22-226. Subsets of T cells in rat mediating lethal graft vs. host disease.
8. Thomas, M.L. and J.R. Green. (1983) Eur. J. Immunol. 13, 855-858. Molecular nature of the W3/25 and MRC OX-8 marker antigens for rat T lymphocytes; comparisons with mouse and human antigens.
9. Whiteland, J.L. et al (1995). Immunohistochemical detection of T cell subsets and other leukocytes in paraffin embedded rat and mouse tissues with monoclonal antibodies. J. Histochem. Cytochem. 43: 313-320.

Laboratory Reagent for Research Use Only

® is a registered trademark of Cedarlane