



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

TECHNICALLY *Speaking*

Place your order with CEDARLANE® or your local distributor.

Please contact CEDARLANE® for lot specific information.

Purified Anti-Rat RT1.B Monoclonal Antibody

CL010AP
CL010AP-2
Lot: 1021

DESCRIPTION:

Cedarlane's anti-rat RT1.B monoclonal antibody recognizes Ia from all rat strains tested. It labels a monomorphic determinant of the rat Ia antigen present on the majority of peripheral B lymphocytes, dendritic cells, some macrophages, and certain epithelial cells (2,3,4) but not peripheral T lymphocytes. (4) This antibody also cross reacts with murine Ia antigens (particularly with k and s haplotypes). Analysis of recombinant strains shows that the determinant maps to the I-A region of the mouse MHC (1,2) and correlates with mouse Ia specificity 17 or 18. Thus, the determinant recognized by CL010AP is polymorphic in the mouse. This antibody may be useful for recognizing the rat Ia antigen and can be coupled to Sepharose 4B for purification of rat Ia antigen by antibody affinity chromatography (1).

This clone is reported to work with frozen and paraffin sections (5).

PRESENTATION:

250 µg (CL010AP) or 500 µg (CL010AP-2) purified Ig buffered in PBS + 0.02% NaN₃.

STORAGE AND RECONSTITUTION:

Store at 4°C. For long term storage, aliquot and freeze unused portion at -20°C in volumes appropriate for single usage. Avoid freeze/thaw cycles.

For more information or to place an order please contact...

CEDARLANE®
LABORATORIES LIMITED
5516 - 8th Line, R.R.#2, Hornby, Ontario, CANADA L0P 1E0



toll free: 1-800-268-5058
in North America

phone: (905) 878-8891 • fax: (905) 878-7800

or visit our website for a list of our international distributors including contact information
website: www.cedarlanelabs.com • e-mail: info@cedarlanelabs.com

SPECIFICATIONS:

CLONE: MRC OX-6

HYBRIDOMA PRODUCTION:**IMMUNIZATION:**

IMMUNOGEN: Rat thymocyte membrane glycoprotein.

IMMUNOCYTE DONOR: BALB/c spleen

FUSION PARTNER: P3-NS1-1-Ag4 (NS1/1)

SPECIFICITY: Rat RT1.B (Rat Ia-non-polymorphic)

Ig CLASS: Mouse IgG₁

PRESENTATION: Purified Ig in PBS with the addition of 0.02% NaN₃.
(Ascites purified via Protein G Chromatography)

ANTIBODY CONCENTRATION: 1.0 mg/ml

FLOW CYTOMETRY ANALYSIS:

1. Prepare cell suspension in media A. For cell preparations, deplete the red blood cell population with Lympholyte ® Rat cell separation medium (CL5040).
2. Wash 2 times.
3. Resuspend cells to 1×10^6 cells in approximately 50 μ l media A in a microcentrifuge tube. (ie. 50 μ l of cells resuspended to 2×10^7 cells / ml).
the contents of 1 tube represent 1 test.
4. To each tube add 50 μ l of a 1:250-1:1000 dilution of **CL010AP or CL010AP-2**.
5. Vortex the tubes to ensure thorough mixing of antibody and cells.
6. Incubate the tubes for 30 minutes at 4°C.
7. Wash 2 times at 4°C.
8. Add 100 μ l of secondary antibody **CLCC30201** (Goat anti-mouse IgG (H+L)-FITC conjugate) at 1:700 dilution.
9. Incubate tubes at 4°C for 30-60 minutes. (It is recommended that the tubes are protected from light since most fluorochromes are light sensitive).
10. Wash 2 times at 4°C in Media B.
11. Resuspend the cell pellet in 50 μ l ice cold Media B.
12. Transfer to suitable tubes for flow cytometric analysis containing 15 μ l of propidium iodide at 0.5 mg/ml in phosphate buffered saline. (This stains dead cells by intercalating DNA).

MEDIA

- A. Phosphate buffered saline (pH 7.2) + 5% normal serum of host species + sodium azide (100 μ l of 2M sodium azide in 100 mls).
- B. Phosphate buffered saline (pH 7.2) + 0.5% bovine serum albumin + sodium azide (100 μ l of 2M sodium azide in 100 mls).

FLOW CYTOMETRIC ANALYSIS

Rat Strain: Wistar Rat

Cell Concentration: 1×10^6 cells per test

Antibody Concentration: 1:400

Isotypic Control: Mouse IgG₁, k


***RECOMMENDED DILUTION FOR USE:** 1:250 -1:1000

CELL SOURCE

Thymus
Spleen
Lymph Node

PERCENT STAINING

8.2%
48.1%
21.8%

**LFL1**

Cell Source: Splenocytes

Percentage Of Cells Stained Above Control: 48.1%

N.B. Appropriate control samples should always be included in any labelling studies.

***For optimal results in various applications, it is recommended that each investigator determine dilutions appropriate for individual use.**

STRAIN DISTRIBUTION:**PROCEDURE:** as above**ANTIBODY CONCENTRATION:** 1:500**STRAINS TESTED:** Wistar, Buffalo, Brown Norway, Fischer , Lewis**POSITIVE:** Wistar, Buffalo, Fischer , Lewis, Brown Norway**NEGATIVE:** none**REFERENCES:**

1. McMaster, W.R. and A.F. Williams. (1979) Identification of Ia glycoproteins in rat thymus and purification from rat spleen. *Eur. J. Immunol.* 9, 426-433.
2. Barclay, A.N. (1981) The localization of populations of lymphocytes defined by monoclonal antibodies in rat lymphoid tissues. *Immunology.* 42, 593-600.
3. Barclay, A.N. and G. Mayrhofer. (1981) Bone Marrow origin of Ia-positive cells in the medulla of rat thymus. *J. Exp. Med.* 153, 1666-1671.
4. Barclay, A.N. (1981) Different reticular elements in rat lymphoid tissue identified by localization of Ia, Thy-1 and MRC OX-2 antigens. *Immunology.* 44, 727-736.
5. Whiteland, J.L et al (1995). Immunohistochemical detection of T cell subsets and other leukocytes in paraffin embedded rat and mouse tissues with monoclonal antibodies .*J. Histochem. Cytochem.* 43: 313-320.

FOR LABORATORY RESEARCH ONLY

(Not to be administered to humans or animals nor used for any drug purpose).

® is a registered trademark of Cedarlane Laboratories Ltd.