



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

TECHNICALLY *Speaking*

Place your order with CEDARLANE® or your local distributor.
Please contact CEDARLANE® for lot specific information.

Purified Anti-Rat CD3 Monoclonal Antibody

CL020AP
LOT: 41000209

DESCRIPTION:

Cedarlane's anti-rat CD3 monoclonal antibody recognizes a rat T cell surface antigen having a molecular weight of 25,000 Da. This surface antigen appears to be associated with a 92,000 Da heterodimer with subunits of 52,000 and 43,000 Da.

The antibody reacts with rat T cells but not B cells. It has been reported that immunohistochemically it stains the lymphocytes in the periarterial lymphatic sheaths of the spleen. It stains the medullary cells of the thymus very strongly and the cortical cells much more weakly.

The antibody has the ability to induce rat T cell proliferation in the presence of PMA or when cross - linked to a solid support.

All of these properties resemble those of CD3 in human and mouse systems (1).

This antibody is suitable for use in flow cytometry.

PRESENTATION:

200 µg purified Ig buffered in PBS and 0.09% sodium azide (NaN₃).

STORAGE/STABILITY:

Store at 4°C. For long term storage, aliquot and freeze unused portion at -20°C in volumes appropriate for single usage. Avoid freeze/thaw cycles. Mix thoroughly after thawing before use. If reagent is to be diluted, it is recommended that only the quantity to be used within one week be diluted.

Continued Overleaf.....

For more information or to place an order please contact...

CEDARLANE®
LABORATORIES LIMITED



toll free: 1-800-268-5058
in North America

phone: (905) 878-8891 • fax: (905) 878-7800

5516 - 8th Line, R.R.#2, Hornby, Ontario, CANADA L0P 1E0

or visit our website for a list of our international distributors including contact information
website: www.cedarlanelabs.com • e-mail: info@cedarlanelabs.com

SPECIFICATIONS:

Clone: 1F4

Specificity: Rat CD3

Ig Class: Mouse IgM

Format: Purified Ig buffered in PBS and 0.09% NaN₃.

Antibody Concentration: 1.0 mg/ml

FLOW CYTOMETRY ANALYSIS:

Method:

1. Prepare a cell suspension in media A. For cell preparations, deplete the red blood cell population with Lympholyte[®]-Rat cell separation medium (CL5040).
2. Wash 2 times.
3. Resuspend the cells to a concentration of 2×10^7 cells/ml in media A. Add 50µl of this suspension to each tube (each tube will then contain 1×10^6 cells, representing 1 test).
4. To each tube, add 1.0 µg* of **CL020AP**.
5. Vortex the tubes to ensure thorough mixing of antibody and cells.
6. Incubate the tubes for 30 minutes at 4°C.
7. Wash 2 times at 4°C.
8. Add 100 µl of secondary antibody (FITC Goat anti-mouse IgM (H+L)) at the recommended dilution.
9. Incubate the tubes at 4°C for 30-60 minutes.
(It is recommended that the tubes are protected from light since most fluorochromes are light sensitive).
10. Wash 2 times at 4°C in media B.
11. Resuspend the cell pellet in 50 µl ice cold media B.
12. Transfer to suitable tubes for flow cytometric analysis containing 15 µl of propidium iodide at 0.5 mg/ml in PBS. This stains dead cells by intercalating in DNA.

Media:

- A. Phosphate buffered saline (pH 7.2) + 5% normal serum of host species + sodium azide (100 µl of 2M sodium azide in 100 mls).
- B. Phosphate buffered saline (pH 7.2) + 0.5% Bovine serum albumin + sodium azide (100 µl of 2M sodium azide in 100 mls).

N.B. Appropriate control samples should always be included in any labelling studies.

*** For optimal results in various applications, it is recommended that each investigator determine dilutions appropriate for individual use.**

REFERENCES:

1. Nicolls, M.G., Aversa, G.A., Pearce, N.W., Spinelli, A., Berger, M.F., Gurley, K.E. and Hall, B.M. 1992 Induction of long term specific tolerance to allografts in vivo by therapy with a monoclonal antibody to rat CD3. *Transplantation* **55**:459-468.
2. Tanaka, T., Masuko, T., Yagita, H., Tamura, T. and Hashimoto, Y. 1989 Characterization of a CD3 - like rat T cell surface antigen recognized by a monoclonal antibody. *J. of Immunol.* **142**:2791-2795.

Laboratory Reagent For Research Use Only

® is a Registered Trademark of Cedarlane Laboratories Limited.