



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Technically  
Speaking

**CEDARLANE**<sup>®</sup>  
[www.cedarlanelabs.com](http://www.cedarlanelabs.com)



Conveniently Delivering You Today's Innovations  
for the Science of Tomorrow™

**Anti-Rat CD25 (IL-2R)  
Monoclonal Antibody**

Catalogue#	Format	Size	Concentration	Isotype Control
CL039A	Ascites	0.5ml	NA	CLCMG100
CL039AP/-2	Purified	250µg/500µg	1.0 mg/ml	CLCMG100
CL039B/-5	Biotin	100µg/500µg	0.1 mg/ml	CLCMG115
CL039F/-5	FITC	100µg/500µg	0.1 mg/ml	CLCMG101
CL039PE/-4	PE	50µg/200µg	0.1 mg/ml	CLCMG104
CL039NA	No Azide	1.0mg	1.0 mg/ml	CLCMG100

Isotype: Mouse IgG<sub>1</sub>

**DESCRIPTION:**

Cedarlane's anti-rat Interleukin-2 receptor monoclonal antibody recognizes the smaller (alpha subunit) 55kD chain of the IL-2 receptor found on activated rat T cells, thymic dendritic cells but not resting lymphocytes (1). CL039 binds to the rat interleukin-2 receptor and has proven to be an important marker for activated T cells (2).

This clone is reported to work with frozen and paraffin sections (3).

**PRESENTATION:**

**Ascites:** Lyophilized.

**Purified:** Purified IgG buffered in PBS and 0.02% NaN<sub>3</sub>. (Purified from ascitic fluid via Protein G Chromatography). For maximum recovery of contents, spin down tube before use.

**Biotin, FITC and PE:** Biotin/FITC/PE conjugated IgG buffered in PBS, 0.02% NaN<sub>3</sub> and EIA grade BSA as a stabilizing protein to bring total protein concentration to 4-5 mg/ml.

**No Azide:** Purified Ig buffered in PBS with no preservative; 0.2µm sterile filtered.

**STORAGE/STABILITY:**

Store **Ascites** at -20°C. For all other formats, store at 4°C. DO NOT FREEZE **PE** conjugates. Handle NA format under aseptic conditions. For long term storage (**Purified, Biotin** and **FITC**), aliquot and freeze unused portions at -20°C in volumes appropriate for single usage. Avoid freeze/thaw cycles.

Visit our website for your local distributor.

**CEDARLANE**<sup>®</sup>



[www.cedarlanelabs.com](http://www.cedarlanelabs.com)

An ISO 9001:2000 and ISO 13485:2003  
registered company.

In CANADA: Toll Free: 1-800-268-5058

4410 Paletta Court, Burlington, ON L7L 5R2 ph: (289) 288-0001, fax: (289) 288-0020  
e-mail: [general@cedarlanelabs.com](mailto:general@cedarlanelabs.com)

In the USA: Toll Free: 1-800-721-1644

1210 Turrentine Street, Burlington, NC 27215 ph: (336) 513-5135, fax: (336) 513-5138  
e-mail: [service@cedarlanelabs.com](mailto:service@cedarlanelabs.com)

## **SPECIFICATIONS:**

Clone: MRC OX-39

### Hybridoma Production:

Immunization: Immunogen: T blasts from a mixed lymphocyte reaction between purified CD4 positive T cells and irradiated spleens.

Donor: BALB/c spleen

Fusion Partner: NSO/1

Specificity: Rat CD25 (IL-2 Receptor)

## **TEST RESULTS:**

### Tissue Distribution by Flow Cytometry Analysis:

Rat Strain: Wistar

Cell Concentration:  $1 \times 10^6$  cells per test

Antibody Concentration Used:  $1.0 \mu\text{g}/10^6$  cells

### Cell Source


T-cell blasts

Thymus

### Percentage of cells stained above control:

95.3%

0.59%



Cell Source: Con-A activated thymocytes (T-cell blasts)

**N.B. Appropriate control samples should always be included in any labeling studies.**

**\* For optimal results in various applications, it is recommended that each investigator determine dilutions appropriate for individual use.**

## **REFERENCES:**

1. Paterson, D.J., Jeffries, W.A., Green, J.R., Brandon, M.R., Corthesy, P., Puklavec, M. and A.F. Williams. (1987) *Mol. Immunol.* 24, 1281-1290. Antigen of Activated Rat T Lymphocytes Including a Molecule of 50,000 Mr Detected Only on CD4 Positive T Blasts.
2. Barclay, A.N. (1981) *Immunology.* 42 593-600 The Localization of populations of lymphocytes defined with monoclonal antibodies in rat lymphoid tissues.
3. Whiteland, J.L et al (1995). Immunohistochemical detection of T cell subsets and other leukocytes in paraffin embedded rat and mouse tissues with monoclonal antibodies .*J. Histochem. Cytochem.* 43: 313-320.

**FOR RESEARCH USE ONLY**