



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

TECHNICALLY *Speaking*

Place your order with CEDARLANE® or your local distributor.

Please contact CEDARLANE® for lot specific information.

Purified Anti-Rat $\gamma\delta$ TCR Monoclonal Antibody

**CL056AP
CL056AP-2
LOT: 5621**

DESCRIPTION:

Cedarlane's anti-rat $\gamma\delta$ T cell receptor monoclonal antibody detects a T cell-specific heterodimeric 48 and 50 kD cell surface protein that is expressed on greater than 90% of CD3⁺ $\alpha\beta$ TCR— rat peripheral T lymphocytes, and identifies a dense network of dendritic cells in the epidermis as $\gamma\delta$ T cells. Immobilized this antibody induces a strong proliferative response in $\gamma\delta$ T cell cultures supplemented with either IL-2 or IL-4.

Applications: immunoprecipitation, flow cytometry, immunohistochemical staining on frozen sections.

PRESENTATION:

250 μ g (CL056AP) or 500 μ g (CL056AP-2) purified Ig buffered in PBS and 0.02% NaN₃.

STORAGE/STABILITY:

Store at 4°C. For long term storage, aliquot and freeze unused portion at -20°C in volumes appropriate for single usage. Avoid repeated freeze/thaw cycles.

For more information or to place an order please contact...

CEDARLANE®
LABORATORIES LIMITED



*toll free: 1-800-268-5058
in North America*

phone: (905) 878-8891 • fax: (905) 878-7800

5516 - 8th Line, R.R.#2, Hornby, Ontario, CANADA L0P 1E0

or visit our website for a list of our international distributors including contact information
website: www.cedarlanelabs.com • e-mail: info@cedarlanelabs.com

SPECIFICATIONS:

Clone: V65

Hybridoma Production:

Immunization: Immunogen: Rat T Blasts
Donor: BALB/c Spleen

Fusion Partner: X63-Ag 8.653

Specificity: Rat $\gamma\delta$ TCR

Ig Class: Mouse IgG₁

Antibody Concentration: 1.0 mg/ml

FLOW CYTOMETRY ANALYSIS:

Method:

1. Prepare a cell suspension in media A. For cell preparations, deplete the red blood cell population with Lympholyte[®]-Rat cell separation medium (CL5040).
2. Wash 2 times.
3. Resuspend the cells to a concentration of 2×10^7 cells/ml in media A. Add 50 μ l of this suspension to each tube (each tube will then contain 1×10^6 cells, representing 1 test).
4. To each tube, add 0.5-1.0 μ g* of **CL056AP or CL056AP-2**.
5. Vortex the tubes to ensure thorough mixing of antibody and cells.
6. Incubate the tubes for 30 minutes at 4°C.
7. Wash 2 times at 4°C.
8. Add 100 μ l of secondary antibody **CLCC30204** (PE-Goat anti-mouse IgG (H+L)) at 1:20 dilution.
9. Incubate the tubes at 4°C for 30-60 minutes.
(It is recommended that the tubes are protected from light since most fluorochromes are light sensitive).
10. Wash 2 times at 4°C in media B.
11. Resuspend the cell pellet in 50 μ l ice cold media B.
12. Transfer to suitable tubes for flow cytometric analysis containing 15 μ l of propidium iodide at 0.5 mg/ml in PBS. This stains dead cells by intercalating in DNA.

Media:

- A. Phosphate buffered saline (pH 7.2) + 5% normal serum of host species + sodium azide (100 µl of 2M sodium azide in 100 mls).
- B. Phosphate buffered saline (pH 7.2) + 0.5% Bovine serum albumin + sodium azide (100 µl of 2M sodium azide in 100 mls).

Results:Tissue Distribution by Flow Cytometry Analysis:

Rat Strain: Fischer

Cell Concentration : 1×10^6 cells per test


Antibody Concentration Used: $0.5 \mu\text{g}/10^6$ cells

Isotypic Control: Mouse IgG₁

Cell SourcePercentage of cells stained above control:

Thymus	4.2%
Splenic T Cells*	5.6%

*(T cells isolated with CL102 - Cedarlane's Rat T Cell Recovery Column Kit)



Cell Source: Splenic T-cells

Percentage of cells stained above control: 5.6%

N.B. Appropriate control samples should always be included in any labelling studies.

*** For optimal results in various applications, it is recommended that each investigator determine dilutions appropriate for individual use.**

REFERENCES:

1. Kühnlein, P., Park, H-J., Herrmann, T., Elbe, A., and Hünig, T. (1994). Identification and Characterization of Rat $\gamma\delta$ T-Lymphocytes in Peripheral Lymphoid Organs, Small Intestine, and Skin With a Monoclonal Antibody to a Constant Determinant of the $\gamma\delta$ T cell Receptor. *J. Immunol.* 153:979-986.

FOR RESEARCH USE ONLY

® is a Registered Trademark of Cedarlane Laboratories Limited.