



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

# TECHNICALLY *Speaking*

Place your order with CEDARLANE® or your local distributor.

*Please contact CEDARLANE® for lot specific information.*

## **Purified Anti-Rat $\alpha\beta$ TCR Monoclonal Antibody (No Azide)**

**CL057NA**  
**LOT: 5721NA**

### **DESCRIPTION:**

Cedarlane's anti-rat  $\alpha\beta$  T cell receptor monoclonal antibody detects a T cell-specific heterodimeric 40 and 46 kD cell surface protein that is expressed on the vast majority, but not all, peripheral rat T cells. CL057NA induces blastoid transformation and IL-2R expression in T-cells when cross linked on an artificial surface. It also induces cell proliferation when exogenous IL-2 is added.

Applications: Immunoprecipitation, flow cytometry, immunohistochemical staining on frozen sections.

### **PRESENTATION:**

1 mg purified Ig buffered in PBS, no preservative; 0.2  $\mu$ m filtered.

### **STORAGE/STABILITY:**

Store at 4°C. For long term storage, aliquot and freeze unused portion at -20°C in volumes appropriate for single usage. Avoid freeze/thaw cycles. Should be handled under aseptic conditions.

For more information or to place an order please contact...

**CEDARLANE®**  
**LABORATORIES LIMITED**



**toll free: 1-800-268-5058**  
**in North America**

phone: (905) 878-8891 • fax: (905) 878-7800

5516 - 8th Line, R.R.#2, Hornby, Ontario, CANADA L0P 1E0

or visit our website for a list of our international distributors including contact information  
website: [www.cedarlanelabs.com](http://www.cedarlanelabs.com) • e-mail: [info@cedarlanelabs.com](mailto:info@cedarlanelabs.com)

**SPECIFICATIONS:**

Clone: R73

**Hybridoma Production:**

Immunization: Immunogen: Rat T Blasts  
Donor: BALB/c Spleen

Fusion Partner: X63-Ag 8.653

Specificity: Rat  $\alpha\beta$  TCR

Ig Class: Mouse IgG<sub>1</sub>

Format: Purified Ig buffered in PBS, no preservative (Purified from BioReactor Supernatant via Protein G Affinity Chromatography.)

Antibody Concentration: 1.0 mg/ml

**FLOW CYTOMETRY ANALYSIS:****Method:**

1. Prepare a cell suspension in media A. For cell preparations, deplete the red blood cell population with Lympholyte®-Rat cell separation medium (CL5040).
2. Wash 2 times.
3. Resuspend the cells to a concentration of  $2 \times 10^7$  cells/ml in media A. Add 50  $\mu$ l of this suspension to each tube (each tube will then contain  $1 \times 10^6$  cells, representing 1 test).
4. To each tube, add 0.2 - 0.5  $\mu$ g\* of **CL057NA**.
5. Vortex the tubes to ensure thorough mixing of antibody and cells.
6. Incubate the tubes for 30 minutes at 4°C.
7. Wash 2 times at 4°C.
8. Add 100  $\mu$ l of secondary antibody **CLCC30201** (FITC Goat anti-mouse IgG (H+L)) at 1:500 dilution.
9. Incubate the tubes at 4°C for 30-60 minutes.  
(It is recommended that the tubes are protected from light since most fluorochromes are light sensitive).
10. Wash 2 times at 4°C in media B.
11. Resuspend the cell pellet in 50  $\mu$ l ice cold media B.
12. Transfer to suitable tubes for flow cytometric analysis containing 15  $\mu$ l of propidium iodide at 0.5 mg/ml in PBS. This stains dead cells by intercalating in DNA.

Media:

- A. Phosphate buffered saline (pH 7.2) + 5% normal serum of host species + sodium azide (100 µl of 2M sodium azide in 100 mls).
- B. Phosphate buffered saline (pH 7.2) + 0.5% Bovine serum albumin + sodium azide (100 µl of 2M sodium azide in 100 mls).

Results:Tissue Distribution by Flow Cytometry Analysis:

Rat Strain: Fischer

Cell Concentration :  $1 \times 10^6$  cells per test

Antibody Concentration Used:  $0.2 \mu\text{g}/10^6$  cells


Isotypic Control: Mouse IgG<sub>1</sub>

Cell SourcePercentage of cells stained above control:

Thymus 61.1%

Splenic T Cells\* 88.8%

\*(T cells isolated with CL102 - Cedarlane's Rat T Cell Recovery Column Kit)



Cell Source: Splenic T-cells

Percentage of cells stained above control: 88.8%

**N.B.** Appropriate control samples should always be included in any labelling studies.

\* For optimal results in various applications, it is recommended that each investigator determine dilutions appropriate for individual use.

**Strain Distribution by Flow Cytometry Analysis:**

Procedure: see page 2

Cell Concentration :  $1 \times 10^6$  cells per test

Antibody Concentration Used:  $0.2 \mu\text{g}/10^6$  cells

Strains Tested: Buffalo, Wistar, Fischer 344, Brown Norway

Positive: Buffalo, Wistar, Fischer 344, Brown Norway

Negative: none

**REFERENCES:**

1. Hunig, T., Wallny, H.J., Hartley, J.K., Lawetzky, A., and Tiefenthaler, G. (1989). A Monoclonal Antibody to a Constant Determinant of the Rat T-Cell Antigen Receptor that Induces T-Cell Activation. *J. Exp. Med.* 169:73-86.

**FOR RESEARCH USE ONLY**

® is a Registered Trademark of Cedarlane Laboratories Limited.