



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

TECHNICALLY *Speaking*

Place your order with CEDARLANE® or your local distributor.

Please contact CEDARLANE® for lot specific information.

PE Anti-Rat CD26 Monoclonal Antibody

CL061PE
CL061PE-4
LOT:

DESCRIPTION:

Cedarlane's Anti-Rat CD26 monoclonal antibody (Clone: OX-61) recognizes and binds to the enzyme dipeptidyl peptidase IV (DPP-IV) present on rat thoracic duct lymphocytes and rat liver cells. The ectoenzyme DPP-IV is a bile canalicular cell surface molecule that has widespread tissue distribution. DPP-IV activity is specifically depleted from solubilized liver homogenates by CD26 bound to Sepharose. Under reducing conditions, the molecular weight of CD26 is 110kD. This antibody is the rat equivalent of human CD26. The N-terminal amino acid sequence of CD26 was sequenced and indicated identity with part of the predicted N-terminus of the bile canalicular molecule GP110. The cloning of DPP-IV permitted a comparison between the molecule recognized by monoclonal antibody CD26, GP110, and DPP-IV. It has been concluded that these three molecules are almost certainly identical.

This product is suitable for use in flow cytometry.

PRESENTATION:

50 µg (CL061PE) or 200 µg (CL061PE-4) PE conjugated IgG buffered in PBS, 0.02% NaN₃, and EIA grade BSA as a stabilizing protein to bring the total protein concentration to 4-5 mg/ml.

STORAGE/STABILITY:

Store at 4°C. **DO NOT FREEZE!** Avoid prolonged exposure to light.

SPECIFICATIONS:

Clone: OX-61

Hybridoma Production:

Immunization:

Immunogen: Thoracic duct lymphocytes from a mesenteric lymphodectomised rat

Donor: BALB/c mouse spleen

Fusion Partner: P3-NSI/1 Ag4-1

Continued Overleaf...

For more information or to place an order please contact...

CEDARLANE®
LABORATORIES LIMITED



toll free: 1-800-268-5058
in North America

phone: (905) 878-8891 • fax: (905) 878-7800

5516 - 8th Line, R.R.#2, Hornby, Ontario, CANADA L0P 1E0

or visit our website for a list of our international distributors including contact information

website: www.cedarlanelabs.com • e-mail: info@cedarlanelabs.com

Specificity: Rat CD26

Ig Class: Mouse IgG2a

Format: R-PE conjugated IgG buffered in PBS, 0.02% NaN₃ and EIA grade BSA as a stabilizing protein to bring the total protein concentration to 4-5 mg/ml.

Antibody Concentration: 0.1 mg/ml

FLOW CYTOMETRY ANALYSIS:

1. Prepare cell suspension in Media A. For cell preparations, deplete the red blood cell population with Lympholyte[®]-R cell separation medium (CL5040).
2. Wash 2 times.
3. Resuspend the cells to a concentration of 2×10^7 cells/ml in Media A. Add 50 μ l of this suspension to each tube (each tube will then contain 1×10^6 cells representing 1 test).
4. To each tube add ~ μ g of **CL061PE or CL061PE-4** per 10^6 cells*.
5. Vortex the tubes to ensure thorough mixing of antibody and cells.
6. Incubate the tubes for 30 minutes at 4°C. (It is recommended that the tubes are protected from light, since most fluorochromes are light sensitive.)
7. Wash 2 times at 4°C.
7. Resuspend the cell pellet in 50 μ l ice cold Media B.
8. Transfer to suitable tubes for flow cytometric analysis containing 15 μ l of propidium iodide at 0.5 mg/ml in phosphate buffered saline. (This stains dead cells by intercalating DNA).

Media:

- A. Phosphate buffered saline (pH 7.2) + 5% normal serum of host species + sodium azide (100 μ l of 2M sodium azide in 100 mls).
- B. Phosphate buffered saline (pH 7.2) + 0.5% Bovine serum albumin + sodium azide (100 μ l of 2M sodium azide in 100 mls).

Results:

Tissue Distribution by Flow Cytometry Analysis:

Rat Strain:

Cell Concentration: 1×10^6 cells per test

Antibody Concentration Used: μ g/ 10^6 cells

Isotypic Control:

Percentage of cells stained above control: %

N.B. Appropriate control samples should always be included in any labeling studies.

*** For optimal results in various applications, it is recommended that each investigator determine dilutions appropriate for individual use.**

REFERENCES:

1. McCaughan, G.W., Wickson, J.E., Creswick, P.F., Gorrell, M.D., 1990. Identification of the Bile Canalicular Cell Surface Molecule CP110 as the Ectopeptidase Dipeptidyl Peptidase IV: An Analysis by Tissue Distribution, Purification and N-Terminal Amino Acid Sequence. *Hepatology*; 11:534

FOR RESEARCH USE ONLY

® is a Registered Trademark of Cedarlane Laboratories Limited.