



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Technically
Speaking

CEDARLANE[®]
www.cedarlanelabs.com



Conveniently Delivering You Today's Innovations
for the Science of Tomorrow™

Anti-Rat Crry Monoclonal Antibody

Catalogue#	Format	Size	Concentration	Isotype Control
CL064AP	Purified	250ug	1.0 mg/ml	CLCMG100
CL064B	Biotin	100ug	0.1 mg/ml	CLCMG115
CL064F	FITC	100ug	0.1 mg/ml	CLCMG101

Isotype: Mouse IgG1

DESCRIPTION:

Cedarlane's anti-Rat Crry monoclonal antibody (Clone: 5I2) is a rat-specific membrane complement regulator that can inhibit both antibody-induced classical pathway and alternative pathway complement activation. Crry plays a more dominant role than DAF in regulating the alternative pathway of complement and in preventing spontaneous complement damage of rat erythrocytes, whereas DAF and Crry are both expressed and are equally effective in preventing antibody-induced runaway complement activation on rat erythrocytes.

This antibody is suitable for use in flow cytometry⁶ and immunohistochemistry (acetone-fixed frozen sections).^{3,4,5,6}

PRESENTATION:

Purified: Purified IgG buffered in PBS and 0.02% NaN₃. (Purified from ascitic fluid via Protein G Chromatography). For maximum recovery of contents, spin down tube before use.

Biotin and FITC: Biotin/FITC conjugated IgG buffered in PBS, 0.02% NaN₃ and EIA grade BSA as a stabilizing protein to bring total protein concentration to 4-5 mg/ml.

STORAGE/STABILITY:

Store at 4°C. For long term storage, aliquot and freeze unused portion at -20°C in volumes appropriate for single usage. Avoid freeze/thaw cycles.

SPECIFICATIONS:

Clone: 5I2

Immunogen: Erythrocytes from a C3 mutated rat

Specificity: Rat Crry

Ig Class: Mouse IgG1

Visit our website for your local distributor.

CEDARLANE[®]



www.cedarlanelabs.com

An ISO 9001:2000 and ISO 13485:2003
registered company.

In CANADA: Toll Free: 1-800-268-5058

4410 Paletta Court, Burlington, ON L7L 5R2 ph: (289) 288-0001, fax: (289) 288-0020
e-mail: general@cedarlanelabs.com

In the USA: Toll Free: 1-800-721-1644

1210 Turrentine Street, Burlington, NC 27215 ph: (336) 513-5135, fax: (336) 513-5138
e-mail: service@cedarlanelabs.com

TEST RESULTS:

Tissue Distribution by Flow Cytometry Analysis:

Rat Strain: Wister

Cell concentration: 1×10^6 cells per test

Antibody Concentration Used: $0.1 \mu\text{g}/10^6$ cells

Cell Source

Bone Marrow

Blood



Spleen

Percentage of cells stained above control

99.6%

55.9 %

59.6%



Wistar rat splenocytes (left) and bone marrow (right) were stained with anti-Crry (clone: 5I2) (filled histogram) or mouse IgG2a isotype control (open histogram).

N.B. Appropriate control samples should always be included in any labeling studies.

*** For optimal results in various applications, it is recommended that each investigator determine dilutions appropriate for individual use.**

REFERENCES:

1. Molina, H., Miwa, *et al.* 2002. Complement-mediated clearance of erythrocytes: mechanism and delineation of the regulatory roles of Crry and DAF. *The American Society of Hematology*
2. Ohta, R., *et al.* 1999. Characteration of Mouse DAF on Trasfectant Cells Using Monoclonal Antibodies Which Recognize Different Epitopes. Nagoya City University School of Medicine and The Scripps Research Institute.
3. Nishikawa, K., *et al.* 1996. Local inflammation caused by a monoclonal antibody that blocks the function of the rat membrane inhibitor of C3 convertase. *The Journal of Immunology*. **156**: 1182-1188.
4. Nomura, A., *et al.* 1995. Tubulointerstitial injury induced in rats by a monoclonal antibody that inhibits function of a membrane inhibitor of complement. *J. Clin. Invest.* Vol. 96. 2348-2356.
5. Matsuo, S., *et al.* 1994. In vivo effects of monoclonal antibodies that functionally inhibit complement regulatory proteins in rats. *J. Exp. Med.* Vol. 180. 1619-1627.
6. Nishikage, H., *et al.* 1995. The role of a complement regulatory protein in rat mesangial glomerulonephritis. *J. Am. Soc. Nephrol.* **6**: 234-242.

FOR RESEARCH USE ONLY

® is a Registered Trademark of Cedarlane