



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Technically  
Speaking

**CEDARLANE**<sup>®</sup>  
[www.cedarlanelabs.com](http://www.cedarlanelabs.com)



Conveniently Delivering You Today's Innovations  
for the Science of Tomorrow™

**Anti-Rat CD200 (OX-2)  
Monoclonal Antibody**

Catalogue#	Format	Size	Concentration	Isotype Control
<b>CL082A</b>	Ascites	0.5 mL	N/A	CLCMG100
<b>CL082AP</b>	Purified	250µg	1.0 mg/ml	CLCMG100
<b>CL082B/-5</b>	Biotin	100µg/500µg	0.1 mg/ml	CLCMG115
<b>CL082F/-5</b>	FITC	100µg/500µg	0.1 mg/ml	CLCMG101
<b>CL082PE/-4</b>	PE	50µg/200µg	0.1 mg/ml	CLCMG104

Isotype: Mouse IgG1

**DESCRIPTION:**

Cedarlane's mouse anti-rat CD200 (OX-2) monoclonal antibody recognizes a monomorphic determinant present on rat thymocytes, brain, follicular dendritic cells in lymphoid organs, vascular endothelium, and at low levels on some smooth muscle and B lymphocytes. The purified brain and thymocyte OX-2 antigens are glycoproteins with apparent M.W. of 41 KDa and 47 KDa respectively. The amino acid composition of brain and thymocyte OX-2 antigen are very similar and are antigenically similar to those found on other tissues. The carbohydrate composition shows that this antigen is highly glycosylated (brain OX-2 - 24% and thymocyte OX-2 - 33% carbohydrate by weight).

The OX-2 antigen shows similarities to the Thy-1 antigen in its odd pattern of tissue distribution, carbohydrate composition and characteristic migration on SDS-PAGE. Also, the OX-2 antigens, like Thy-1 antigens, have homologies with immunoglobulin domains; the overall structure of OX-2 is similar to an Ig light chain or the T cell receptor  $\beta$  chain. Because of its distribution, it is thought to play a role in mediating recognition events at cell surfaces.

This clone can be used in flow cytometry, affinity chromatography, immunohistochemistry and in binding assays. It is also useful for labeling the follicular dendritic cells thought to be involved in the generation of B cell memory as it does not label the Ia-positive dendritic cells present in the T-dependent areas of lymphoid organs.

**PRESENTATION:**

**Ascites:** Lyophilized.

**Purified:** Purified IgG buffered in PBS and 0.02% NaN<sub>3</sub>. (Purified from ascitic fluid via Protein G Chromatography). For maximum recovery of contents, spin down tube before use.

**Biotin, FITC and PE:** Biotin/FITC/PE conjugated IgG buffered in PBS, 0.02% NaN<sub>3</sub> and EIA grade BSA as a stabilizing protein to bring total protein concentration to 4-5 mg/ml.

Visit our website for your local distributor.

**CEDARLANE**<sup>®</sup>



[www.cedarlanelabs.com](http://www.cedarlanelabs.com)

An ISO 9001:2000 and ISO 13485:2003  
registered company.

In CANADA: Toll Free: 1-800-268-5058

4410 Paletta Court, Burlington, ON L7L 5R2 ph: (289) 288-0001, fax: (289) 288-0020  
e-mail: [general@cedarlanelabs.com](mailto:general@cedarlanelabs.com)

In the USA: Toll Free: 1-800-721-1644

1210 Turrentine Street, Burlington, NC 27215 ph: (336) 513-5135, fax: (336) 513-5138  
e-mail: [service@cedarlanelabs.com](mailto:service@cedarlanelabs.com)

### **STORAGE/STABILITY:**

Store **Ascites** at -20°C. For all other formats, store at 4°C. **DO NOT FREEZE PE** conjugates. For long term storage (**Purified, Biotin** and **FITC**), aliquot and freeze unused portion at -20°C in volumes appropriate for single usage. Avoid freeze/thaw cycles.

### **SPECIFICATIONS:**

Clone: OX-2

Hybridoma Production: Spleen cells from BALB/c mice immunized with rat thymocyte membrane glycoproteins were fused with mouse NS-1 myeloma cells.

Specificity: Rat thymocytes, brain, follicular dendritic cells in lymphoid organs, vascular endothelium, and at low levels on some smooth muscle and B lymphocytes

### **TEST RESULTS:**


#### Tissue Distribution by Flow Cytometry Analysis:

Rat Strain: Wistar

Cell Concentration:  $1 \times 10^6$  cells per test

Antibody Concentration Used:  $0.5 \mu\text{g}/10^6$  cells

<u>Cell Source</u>	<u>Percentage of cells stained above control:</u>
Thymus	98.7%
Spleen	35.7%
Lymph Node	39.9%



**N.B. Appropriate control samples should always be included in any labeling studies.**

**\* For optimal results in various applications, it is recommended that each investigator determine dilutions appropriate for individual use.**

### **REFERENCES:**

- 1) McMaster, Robert W. and Williams, Alan F. (1979) *Eur. J. Immunol.* 9:426-433. Identification of Ia glycoproteins in rat thymus and purification from rat spleen.
- 2) Barclay, A. Neil. (1981) *Immunology* 44:727-736. Different reticular elements in rat lymphoid tissues identified by localization of Ia, Thy-1 and MRC OX-2 antigens.
- 3) Barclay, A. Neil and Ward, Harry A. (1982) *Eur. J. Immunol.* 129:447-458. Purification and Chemical Characterization of Membrane Glycoproteins from Rat Thymocytes and Brain, Recognized by Monoclonal Antibody MRC OX-2.
- 4) Clark, Melanie J., Gagnon, Jean, Williams, Alan F. and Barclay, A. Neil. (1985) *EMBO Journal* 4:113-118. MRC OX-2 antigen: a lymphoid/neuronal membrane glycoprotein with a structure like a single immunoglobulin light chain.
- 5) Barclay, A. Neil. (1981) *Immunology* 42:593-600. The localization of populations of lymphocytes defined by monoclonal antibodies in rat lymphoid tissues.

**FOR RESEARCH USE ONLY**  
® is a registered trademark of Cedarlane