



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

TECHNICALLY Speaking

Place your order with CEDARLANE® or your local distributor.

Please contact CEDARLANE® for lot specific information.

PE Mouse Anti-Human p75 NGFR (low affinity receptor)

CL10013PE

Lot:

Description: PE conjugated Mouse anti-Neuronal Growth Factor Receptor (NGFR), p75, low affinity receptor. Detects a protein with a molecular weight of 75 kDa.

Antibody Content: 100 µg

Clone: ME20.4

Immunogen: WM245 melanoma cells

Isotype: Mouse IgG1, k

Formulation: Purified via protein A in 1.0 ml PBS (20mM), 0.14M NaCl, pH 7.3. This product contains 0.1% sodium azide (NaN₃) and 0.2% gelatin.

Applications: Suitable for use in flow cytometry.

Flow Cytometry Protocol:

1. Add 10 µl of antibody to 1 x 10⁶ cells.
2. Incubate 30 minutes on ice in PBS containing 2-5% BSA.
3. Wash via centrifugation.
4. Analyze by flow cytometry.

Positive Control Cell Line: HS294T from ATCC

Procedure For General Staining Using Flow Cytometry: (For Non-Adherent Cells):

1. Add 0.3-1.0 µg anti-NGFR FITC or PE in 10 µl to one million cells in 100 µl PBS, 2% BSA.
2. Incubate on ice for 30 minutes.
3. Add 1.0 ml PBS, BSA and centrifuge for 5 minutes at 500xg to wash cells.
4. Suction off PBS, BSA and add 1.0 ml fresh PBS, BSA.
5. Analyze by flow cytometry.
6. HS294 T-cell line from ATCC can be used for positive control.

References:

1. Ross, et al., Characterization of nerve growth factor receptor in neural crest tumors using monoclonal antibodies, Proc. Natl. Acad. Sci. 81:6681-6685, 1985.

Continued overleaf...

For more information or to place an order please contact...

CEDARLANE®
LABORATORIES LIMITED



toll free: 1-800-268-5058
in North America

phone: (905) 878-8891 • fax: (905) 878-7800

5516 - 8th Line, R.R.#2, Hornby, Ontario, CANADA L0P 1E0

or visit our website for a list of our international distributors including contact information
website: www.cedarlanelabs.com • e-mail: info@cedarlanelabs.com

2. Valtieri, et al., Efficient transfer of Selectable and membrane reporter genes in hematopoietic progenitor and stem cells purified from human peripheral blood., *Cancer Research* 54: 4398-4404, 1994.
3. Stove, et al. Human Immunodeficiency Virus Nef Induces Rapid Internalization of the T-Cell Coreceptor CD8 . *Journal of Virology*, 79 (17):11422-11433, 2005.
4. Vissavajhala, et al. Structural domains of the extracellular domain of human nerve growth factor receptor detected by partial proteolysis. *Arch. Biochem. Biophys.* 294(1): 244-252, 1992.

FOR RESEARCH USE ONLY

JCr 3/20/06