



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Technically
Speaking

CEDARLANE[®]
www.cedarlanelabs.com



Conveniently Delivering You Today's Innovations
for the Science of Tomorrow™

**Anti-Human CDCrel-1/Sept5
Monoclonal Antibody**

Catalogue#	Format	Size	Concentration	Isotype Control
CL2708AP	Purified	200 µg	1.0 mg/ml	CLCMG100
CL2708B	Biotin	100 µg	0.1 mg/ml	CLCMG115
CL2708F	FITC	100 µg	0.1 mg/ml	CLCMG101

Isotype: Mouse IgG1

DESCRIPTION:

CDCrel-1/Septin 5 (Sept5) belongs to the Septin family of nucleotide binding GTPases. Septins were originally described in yeast as cell division cycle regulatory proteins involved in cytokinesis and the regulation of cytoskeletal organization. Sept5 is expressed in cells of the nervous system and is seen to associate primarily with vesicles and membranes through its interaction with the SNARE domain of syntaxin 1A. Through its interaction with syntaxin 1A Sept5 acts to inhibit exocytosis, possibly by regulating vesicle targeting and/or fusion. Recently it was shown that Sept5 is phosphorylated by cyclin-dependent kinase 5 (Cdk5) – p35 decreasing it's binding to syntaxin 1A. This suggests that Cdk5 can modulate synaptic vesicle release by regulating the interactions between Sept5 and syntaxin 1A.

PRESENTATION:

Purified: Purified IgG buffered in PBS and 0.02% NaN₃. (Purified from ascitic fluid via Protein G Chromatography). For maximal recovery of contents, please quick-spin vial before opening.

Biotin and FITC: Biotin/FITC conjugated IgG buffered in PBS, 0.02% NaN₃ and EIA grade BSA as a stabilizing protein to bring total protein concentration to 4-5 mg/mL.

STORAGE/STABILITY:

Store at +4°C. For long term storage, aliquot and freeze unused portion at -20°C in volumes appropriate for single use. Avoid freeze thaw cycles.

APPLICATIONS:

This antibody is suitable for use in Western Blot, ELISA, Flow Cytometry and Immunofluorescence (1:500 dilution).

Continued Overleaf...

Visit our website for your local distributor.

CEDARLANE[®]



www.cedarlanelabs.com

An ISO 9001:2000 and ISO 13485:2003
registered company.

In CANADA: Toll Free: 1-800-268-5058

4410 Paletta Court, Burlington, ON L7L 5R2 ph: (289) 288-0001, fax: (289) 288-0020
e-mail: general@cedarlanelabs.com

In the USA: Toll Free: 1-800-721-1644

1210 Turrentine Street, Burlington, NC 27215 ph: (336) 513-5135, fax: (336) 513-5138
e-mail: service@cedarlanelabs.com

SPECIFICATIONS:

Clone: SP18

Hybridoma Production:

Immunization: Mouse monoclonal antibodies raised against Immunoprecipitate of human brain.

Specificity: This antibody is specific for human CDCrel-1/SEPT5. [also reacts with mouse and rat].

Appropriate control samples should always be included in any labeling studies.

*** For optimal results in various applications, it is recommended that each investigator determine dilutions appropriate for individual use.**

REFERENCES:

1. Beites et al. **The septin Sept5/CDCrel-1 competes with α -SNAP for binding to the SNARE complex.** Biochem. J. 385, 347-353, 2005.
2. Honer, WG et al. **Cingulate cortex synaptic terminal proteins and neural cell adhesion molecule in schizophrenia.** Neuroscience. 78(1), 99-110, 1997.
3. Honer, WG et al. **Human synaptic proteins with a heterogeneous distribution in cerebellum and visual cortex.** Brain Res. 609(1-2), 9-20, 1993.
4. Honer, WG et al. **Monoclonal antibodies to study the brain in schizophrenia.** Brain Res. 500(1-2), 379-83, 1989.
5. Lang, T. and Jahn R. **Core proteins of the secretory machinery.** Handb. Exp. Pharmacol. (184): 107-27, 2008.
6. Taniguchi, M et al. **Phosphorylation of adult type Sept5 (CDCrel-1) by cyclin- dependent kinase 5 inhibits interaction with syntaxin-1.** J. Biol. Chem. 282(11): 7869-76. 2007.
7. Ungermann, C. and Langosch, D. **Functions of SNAREs in intracellular membrane fusion and lipid bilayer mixing.** J. Cell Science. 118, 3819-3828, 2005.

FOR RESEARCH USE ONLY
® is a registered trademark of Cedarlane