



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Technically
Speaking

CEDARLANE[®]
www.cedarlanelabs.com



Conveniently Delivering You Today's Innovations
for the Science of Tomorrow™

Anti-Human Calgranulin A Monoclonal Antibody

Catalogue#	Format	Size	Concentration	Isotype Control
CL2711A	Ascites	0.5 ml	N/A	CLCMG2B00
CL2711AP	Purified	200 µg	1.0 mg/ml	CLCMG2B00
CL2711B	Biotin	100 µg	0.1 mg/ml	CLCMG2B15
CL2711F	FITC	100 µg	0.1 mg/ml	CLCMG2B01

Isotype: Mouse IgG2b

DESCRIPTION:

Cedarlane's Anti-Human Calgranulin A monoclonal antibody reacts with the human Calgranulin A protein. Calgranulin A, also known as MRP-8 and S100A8 is a member of the S100 family of proteins containing 2 EF hand (alpha helix, turn, alpha helix structure) calcium binding motifs. S100 proteins are localized in the cytoplasm and/or nucleus of a wide range of cells and are involved in the regulation of a number of cellular processes such as cell cycle progression and differentiation.

MRP-8 (S100A8) forms a heterodimeric complex with Calgranulin B, (MRP-14, S100A9) in the cytosol of monocyte and neutrophil cell types circulating in peripheral blood. Calgranulin A is found in elevated levels in the serum of cystic fibrosis cases and is also expressed in the skin of patients with psoriasis, eczematous dermatitis and squamous cell carcinoma. We also sell an ascites purified anti-human Calgranulin B monoclonal antibody, CL2712AP.

Reported applications of this antibody include flow cytometry, IF, IHC and Western blots.

PRESENTATION:

Ascites: From ascitic fluid.

Purified: Purified IgG buffered in PBS and 0.02% NaN₃. (Purified from ascitic fluid via Protein G Chromatography). For maximal recovery of contents, please quick-spin vial before opening.

Biotin, FITC: Biotin/FITC/PE conjugated IgG buffered in PBS, 0.02% NaN₃ and EIA grade BSA as a stabilizing protein to bring total protein concentration to 4-5 mg/mL.

STORAGE/STABILITY:

For all formats, store at 4°C. For long term storage (**Ascites, Purified, Biotin, FITC**), aliquot and freeze unused portion at -20°C in volumes appropriate for single usage. Avoid freeze/thaw cycles

Visit our website for your local distributor.

CEDARLANE[®]



www.cedarlanelabs.com

An ISO 9001:2000 and ISO 13485:2003
registered company.

In CANADA: **Toll Free: 1-800-268-5058**

4410 Paletta Court, Burlington, ON L7L 5R2 ph: (289) 288-0001, fax: (289) 288-0020
e-mail: general@cedarlanelabs.com

In the USA: **Toll Free: 1-800-721-1644**

1210 Turrentine Street, Burlington, NC 27215 ph: (336) 513-5135, fax: (336) 513-5138
e-mail: service@cedarlanelabs.com

SPECIFICATIONS:

Clone: CF-145

Hybridoma Production:

Immunization:

Immunogen: Purified granulocyte antigen of human origin

Donor: human leukocytes

Fusion Partner: NSO Mouse Myeloma Partner

Specificity: Binds to human Calgranulin A


TEST RESULTS:

Tissue Distribution by Flow Cytometry Analysis:

Cell Concentration : 1×10^6 cells per tests

Antibody Concentration Used: $0.5 \mu\text{g}/10^6$ cells

Isotypic Control (shaded): FITC Mouse IgG2b (CLCMG2B01)



Human whole blood (granulocyte population) stained with anti-human calgranulin A (clone: CF-145) (filled histogram) or mouse IgG2b isotype control (open histogram).

N.B. Appropriate control samples should always be included in any labeling studies.

*** For optimal results in various applications, it is recommended that each investigator determine dilutions appropriate for individual use.**

REFERENCES:

1. Klein et al. 1996. Identification and Functional Separation of Retinoic Acid Receptor Neutral Antagonists and Inverse Agonists. The Journal of Biological Chemistry Sept; 271(37):22692-22696.
2. Nagpal et al. 1996. Negative Regulation of Two Hyperproliferative Keratinocyte Differentiation Markers by a Retinoic Acid Receptor-specific Retinoid: Insight into the Mechanism of Retinoid Action in Psoriasis. Cell Growth and Differentiation Dec; 7:1783-1791.
3. Thacher et al. 1999. Cell Type and Gene-specific Activity of the Retinoid Inverse Agonist AGN 193109: Divergent Effects from Agonist at Retinoic Acid Receptor γ in Human Keratinocytes. Cell Growth and Differentiation Apr; 10:255-262.

FOR RESEARCH USE ONLY

® is a registered trademark of Cedarlane Laboratories Ltd.