



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Technically
Speaking

CEDARLANE[®]
www.cedarlanelabs.com



Conveniently Delivering You Today's Innovations
for the Science of Tomorrow™

Anti-Human IgG Fd Monoclonal Antibody

Catalogue#	Format	Size	Concentration	Isotype Control
CL6045AP	Purified	200 µg	1.0 mg/ml	CLCMG100
CL6045B	Biotin	100 µg	0.1 mg/ml	CLCMG115
CL6045F	FITC	100 µg	0.1 mg/ml	CLCMG101
CL6045PE	PE	50 µg	0.1 mg/ml	CLCMG104
CL6045HP	HRPO	100 µg	1.0 mg/ml	CLCMG107

Isotype: Mouse IgG₁

DESCRIPTION:

The Fd region of human Immunoglobulins is located at the N-terminal part of the heavy chain and is an essential component of the antigen-binding fragment Fab. Fd, like the light chain, contains a C-terminal constant (C_H1) and N-terminal variable (V_H) domain. The hypervariable regions in both the light chain and Fd determine the specificity of the Fab.

PRESENTATION:

Purified: Purified IgG buffered in PBS and 0.02% NaN₃. (Purified from ascitic fluid via Protein G Chromatography). For maximum recovery of contents, spin down tube before use.

Biotin, FITC, and PE: Biotin/FITC/PE conjugated IgG buffered in PBS, 0.02% NaN₃ and EIA grade BSA as a stabilizing protein to bring total protein concentration to 4-5 mg/ml.

HRPO: HRPO conjugated IgG buffered in PBS with 40 % glycerol (v/v) and EIA grade BSA as a stabilizing protein to bring total protein concentration to 4-5 mg/ml. No NaN₃ with other preservatives.

STORAGE/STABILITY:

For all formats, store at + 4°C. DO NOT FREEZE PE conjugates. For long term storage (**Purified/Biotin/FITC/HRPO**), aliquot and freeze unused portion at -20°C in volumes appropriate for single usage. Avoid freeze/thaw cycles. For Maximal recovery of contents, please quick-spin vial before opening.

APPLICATION:

This antibody is suitable for use in ELISA (both capture and detection antibody), Radioimmunoassay and WB.

* For optimal results in various applications, it is recommended that each investigator determine dilutions appropriate for individual use.

Continued Overleaf.....

Visit our website for your local distributor.

CEDARLANE[®]



www.cedarlanelabs.com

An ISO 9001:2000 and ISO 13485:2003
registered company.

In CANADA: Toll Free: 1-800-268-5058

4410 Paletta Court, Burlington, ON L7L 5R2 ph: (289) 288-0001, fax: (289) 288-0020
e-mail: general@cedarlanelabs.com

In the USA: Toll Free: 1-800-721-1644

1210 Turrentine Street, Burlington, NC 27215 ph: (336) 513-5135, fax: (336) 513-5138
e-mail: service@cedarlanelabs.com

SPECIFICATIONS:

Clone: HP6045

Immunogen: Human IgG Fd

Specificity: Human IgG Fd

ELISA Specificity: human IgG subclass 1 100%
 human IgG subclass 2 100%
 human IgG subclass 3 100%
 human IgG subclass 4 99%
 human IgG Pan (pooled IgG1-4) 100%
 human IgG Fc 0.04%
 human IgG Fab 100%

REFERENCES:

1. Hamilton, R.G. et al; Isoelectric focusing-affinity immunoblot analysis of mouse monoclonal antibodies to the four human IgG subclasses. *Electrophoresis*, 8:127-34, 1987.
2. Papadea, C. et al; Monoclonal antibody-based solid-phase immunoenzymometric assays for quantifying human immunoglobulin G and its subclasses in serum. *Clin. Chem.*, 31: 1940-5, 1985.
3. Jeffries, R. et al; Evaluation of monoclonal antibodies having specificity for human IgG sub-classes: results of an IUIS/WHO collaborative study; [Immunol Lett](#). 10(3-4):223-52, 1985.
4. Reimer, C.B. et al; Evaluation of thirty-one mouse monoclonal antibodies to human IgG epitopes. *Hybridoma*, 3: 263-75, 1984.