



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Technically
Speaking

CEDARLANE[®] 
www.cedarlanelabs.com

Conveniently Delivering You Today's Innovations
for the Science of Tomorrow™

Anti-Human Growth Associated Protein-43 (GAP-43) Polyclonal Antibody

CL7794AP
CL7794AP-S
Lot: 189421

DESCRIPTION:

GAP43 (also known as neuromodulin) is considered a "growth" or "plasticity" protein because it is expressed at high levels in neuronal growth cones during development and axonal regeneration, and it is phosphorylated after long-term potentiation and after learning. It is a nervous tissue-specific protein that localizes to the cytoplasm of neurons and attaches to the membrane via a dual palmitoylation sequence on cysteines 3 and 4. GAP-43 is a major protein kinase C (PKC) substrate.

A null mutation of GAP-43 leads to death within days after birth, due to axon pathfinding defects. Additionally, aberrant expression of GAP43 has been seen in patients diagnosed with schizophrenia and Alzheimer's disease.

PRESENTATION:

100 µg (**CL7794AP**) or 20 µg (**CL7794AP-S**) purified IgG (1 mg/ml) buffered in PBS and 0.02% NaN₃. (Purified from serum via Affinity Chromatography). For maximum recovery of contents, spin down tube before use.

STORAGE/STABILITY:

Store at + 4°C. For long term storage, aliquot and freeze unused portion at -20°C in volumes appropriate for single usage. Avoid freeze/thaw cycles.

SPECIFICATION:

Immunogen: 15 amino acid synthetic peptide located near the N-terminus of human GAP-43.

Specificity: This antibody is specific for human and mouse GAP-43.

IgG Class: Rabbit IgG

Application: This antibody is suitable for use in Western Blot (1 – 3 µg/mL). This antibody has not been tested in other applications.

Continued....

Visit our website for your local distributor.

CEDARLANE[®] 

www.cedarlanelabs.com

An ISO 9001:2015 and ISO 13485:2016
registered company.

In CANADA: Toll Free: 1-800-268-5058


4410 Paletta Court, Burlington, ON L7L 5R2 ph: (289) 288-0001, fax: (289) 288-0020
e-mail: general@cedarlanelabs.com

In the USA: Toll Free: 1-800-721-1644

1210 Turrentine Street, Burlington, NC 27215 ph: (336) 513-5135, fax: (336) 513-5138
e-mail: service@cedarlanelabs.com

TEST RESULTS:

Western Blot:



CL7794AP (1 μ g/ml) staining of human brain (cerebellum) (A) and mouse brain (B) lysates (35 μ g protein in RIPA buffer). Detected by chemiluminescence.

N.B. Appropriate control samples should always be included in any labeling studies.

* For optimal results in various applications, it is recommended that each investigator determine dilutions appropriate for individual use.

REFERENCES:

1. Rosskothén-Kuhl N, Illing RB (2014). Gap43 Transcription Modulation in the Adult Brain Depends on Sensory Activity and Synaptic Cooperation. PLoS ONE. 9 (3): e92624.
2. Mackie Ogilvie C, Rooney SC, Hodgson SV, Berry AC (1998). Deletion of chromosome 3q proximal region gives rise to a variable phenotype. Clinical Genetics. 53 (3): 220–2.

For Research Use Only