



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

# TECHNICALLY *Speaking*

Place your order with CEDARLANE® or your local distributor.

*Please contact CEDARLANE® for lot specific information.*

## **Purified Anti-Mouse I-A<sup>b</sup> Monoclonal Antibody – No Azide**

**CL8702NA**

**LOT: 8222NA**

### **DESCRIPTION:**

Cedarlane's anti- I-A<sup>b</sup> monoclonal antibody reacts with the I-A<sup>b</sup> encoded MHC class II antigen expressed on mouse strains of the H-2<sup>b</sup> haplotype. It also reacts with the I-A<sup>d</sup> encoded MHC class II antigen expressed on mouse strains of the H-2<sup>d</sup> haplotype. The reaction pattern of this antibody with a panel of inbred and recombinant haplotypes demonstrates that the antibody detects the public Ia.m8 determinant. Class II antigens are most highly expressed on antigen-presenting cells including B cells, macrophages, dendritic cells and certain epithelial cells.

This antibody is suitable for use in flow cytometry and can be used to quantitate or eliminate cells bearing the I-A<sup>b</sup> antigen from the appropriate strains of mice.

### **PRESENTATION:**

1.0 mg purified Ig buffered in PBS, no preservative, 0.2µm sterile filtered.

### **STORAGE/STABILITY:**

Store at 4°C. For long term storage, aliquot and freeze unused portion at -20°C in volumes appropriate for single usage. Avoid freeze/thaw cycles. Should be handled under aseptic conditions.

For more information or to place an order please contact...

**CEDARLANE®**  
**LABORATORIES LIMITED**



**toll free: 1-800-268-5058**  
**in North America**

**phone: (905) 878-8891 • fax: (905) 878-7800**

5516 - 8th Line, R.R.#2, Hornby, Ontario, CANADA L0P 1E0

or visit our website for a list of our international distributors including contact information

**website: [www.cedarlanelabs.com](http://www.cedarlanelabs.com) • e-mail: [info@cedarlanelabs.com](mailto:info@cedarlanelabs.com)**

**SPECIFICATIONS:**

Clone: 28-16-8s

Hybridoma Production:

Immunization: Immunogen:C3H.SW spleen  
Donor: C3H

Fusion Partner: SP2/0Ag.14

Specificity: Mouse I-A<sup>b</sup> (Ia.m8)

Ig Class: Mouse IgM

Format: Purified Ig buffered in PBS,no preservative, 0.2µm sterile filtered.  
(Purified from ascitic material via affinity chromatography)

Antibody Concentration: 1.0mg/ml

**CYTOTOXICITY ANALYSIS:**

Method:

1. Prepare a cell suspension from the appropriate tissue in Cedarlane Cytotoxicity Medium<sup>a</sup> (CL95100) or equivalent. Remove red cells and dead cells (where necessary) by purification of viable lymphocytes on Cedarlane Lympholyte<sup>®</sup>-M<sup>b</sup> (CL5030) cell separation medium. After washing, adjust the cell concentration to  $1.1 \times 10^6$  cells per ml in Cytotoxicity Medium.
2. Add the antibody to a final concentration of 1:640 and mix.
3. Incubate for 60 minutes at 4°C.
4. Centrifuge to pellet the cells and discard the supernatant.
5. Resuspend to the original volume in Cedarlane Low-Tox<sup>®</sup>-M Rabbit Complement<sup>c</sup> (CL3051) diluted to the recommended concentration in Cytotoxicity Medium.
6. Incubate for 60 minutes at 37°C.
7. Place on ice.
8. Add Trypan Blue, 10% by volume of 1% Trypan Blue (w/v) added 3-5 minutes before scoring works well. Score live versus dead cells in a hemacytometer. Cytotoxic Index (C. I.) can be calculated as follows:

$$C.I. = 100 \times \frac{\% \text{ cyt (antibody + complement)} - \% \text{ cyt (complement alone)}}{100\% - \% \text{ cyt (complement alone)}}$$

Results:Antibody Titration by Cytotoxicity Analysis:

Cell Source: Splenocytes

Donor: BALB/c


Cell Concentration:  $1.1 \times 10^6$  cells/ml

Complement: Cedarlane Low-Tox® –M Rabbit Complement (CL3051)

Complement Concentration: 1:10

Procedure: Two-stage cytotoxicity as described on page 2

$$\text{C.I.} = 100 \times \frac{\% \text{ cyt (antibody + complement)} - \% \text{ cyt (complement alone)}}{100\% - \% \text{ cyt (complement alone)}}$$

Tissue Distribution by Cytotoxicity Analysis:

Procedure: see page 2

Antibody Concentration Used: 1:640

Strain: BALB/c

<u>Cell Source</u>	<u>C.I.</u>
Thymus	0
Spleen	61
Lymph Node	31
Bone Marrow	17

Strain Distribution by Cytotoxicity Analysis:

Procedure: As above

Antibody Concentration Used: 1:40

<u>Strains</u>	<u>H-2 Loci Alleles</u>									
	<u>K</u>	<u>A<sub>β</sub></u>	<u>A<sub>α</sub></u>	<u>E<sub>β</sub></u>	<u>E<sub>α</sub></u>	<u>C4</u>	<u>C4S</u>	<u>D</u>	<u>+/-</u>	
A.TL	s	k	k	k	k	k		k	d	-
BALB/c	d	d	d	d	d	d		d	d	+
C57BL/6	b	b	b	b	b	b		b	b	+
C3H/He	k	k	k	k	k	k		k	k	-

**CYTOTOXICITY DEPLETION ASSAY:**Method:

1. Prepare a cell suspension from the appropriate tissue in Cedarlane Cytotoxicity Medium<sup>a</sup> (CL95100) or equivalent. Remove red cells and dead cells (where necessary) by purification of viable lymphocytes on Cedarlane Lympholyte<sup>®</sup>-M cell separation medium (CL5030). After washing, adjust the cell concentration to  $1 \times 10^7$  cells per ml in Cytotoxicity Medium.
2. Add the antibody to a final concentration of 1:640 and mix. Alternatively, pellet the cells and resuspend in antibody diluted 1:640 in Cytotoxicity Medium.
3. Incubate for 60 minutes at 4°C.
4. Centrifuge to pellet the cells and discard the supernatant.
5. Resuspend to the original volume in Cedarlane Low-Tox-M<sup>®</sup> Rabbit Complement<sup>c</sup> (CL3051), diluted to the appropriate concentration in Cytotoxicity Medium. (Recommended concentration included with each batch of Low-Tox-M<sup>®</sup> Rabbit Complement.)
6. Incubate for 60 minutes at 37°C.
7. Monitor for percent cytotoxicity at this stage, before further processing. For this purpose, remove a small sample from each tube, dilute 1:10 with medium, and add 1/10 volume of 1% Trypan Blue. After 3-5 minutes, score live versus dead cells in a hemacytometer.
8. For functional studies, remove the dead cells from the treated groups before further processing, particularly if the treated cells are to be cultured. This can be done by layering the cell suspension over a separation medium and centrifuging at room temperature as per the instructions provided. Live cells will form a layer at the interface, while the dead cells pellet. The interface can then be collected and washed in Cytotoxicity Medium before being resuspended in the appropriate medium for further processing. Alternatively, the cells can be washed and resuspended in the appropriate medium for further processing immediately after Step #6, provided that the dead cells will not interfere with subsequent assays.

**NOTES:**

- a. Cedarlane Cytotoxicity Medium (CL95100) is RPMI-1640 with 25 mM Hepes buffer and 0.3% bovine serum albumin (BSA). BSA is substituted for the conventionally used fetal calf serum (FCS) because we have found that many batches of FCS contain complement-dependent cytotoxins to mouse lymphocytes, thus increasing the background killing in the presence of complement. Some batches of BSA also contain complement-dependent cytotoxins, resulting in the same problem. We screen for batches of BSA giving low background in the presence of complement and use the selected BSA for preparing Cedarlane Cytotoxicity Medium.
- b. Cedarlane Lympholyte<sup>®</sup>-M cell separation medium (CL5030) is a density separation medium designed specifically for the isolation of viable mouse lymphocytes. This separation medium provides a high and non-selective recovery of viable mouse lymphocytes, removing red cells and dead cells. The density of this medium is 1.087-1.088. Isolation of mouse lymphocytes on cell separation medium of density 1.077 will result in high and selective loss of lymphocytes and should be avoided.
- c. Rabbit serum provides the most potent source of complement for use with antibodies to mouse cell surface antigens. However, rabbit serum itself is very toxic to mouse lymphocytes. Cedarlane Low-Tox<sup>®</sup>-M Rabbit Complement (CL3051) is absorbed to remove toxicity to mouse lymphocytes, while maintaining its high complement activity. When used in conjunction with Cedarlane Cytotoxicity Medium, this reagent provides a highly potent source of complement with minimal background toxicity.

**FOR RESEARCH USE ONLY**

® is a Registered Trademark of Cedarlane Laboratories Limited.