



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Technically  
Speaking

**CEDARLANE**<sup>®</sup>  
[www.cedarlanelabs.com](http://www.cedarlanelabs.com)



Conveniently Delivering You Today's Innovations  
for the Science of Tomorrow™

**Anti-Mouse CD276 (B7-H3)  
Monoclonal Antibody**

Catalogue#	Format	Size	Concentration	Isotype Control
CL8776AP	Purified	250 µg	1.0 mg/ml	CLCR100
CL8776LE	Low Endotoxin	500 µg	1.0 mg/ml	CLCR100
CL8776B	Biotin	100 µg	0.1 mg/ml	CLCR115
CL8776F	FITC	100 µg	0.1 mg/ml	CLCR101
CL8776PE	PE	50 µg	0.1 mg/ml	CLCR104
CL8776APC	APC	100 µg	0.1 mg/ml	CLCR105

Isotype: Rat IgG<sub>1</sub>, k

**DESCRIPTION:**

CD276 (B7-H3) is a type 1 transmembrane glycoprotein that belongs to the B7 family of co-stimulatory molecules. It's primary expressed at low levels on antigen presenting cells including B cells, macrophages, and dendritic cells. CD276 is also seen to be expressed on various human and murine tumor cells, nasal and airway epithelial cells. CD276 modulates T-cell-mediated immune responses and the development of acute and chronic transplant rejection. Its seen to play a positive regulatory role in bone formation and has a dual role in the bone-immune interface. Additionally, CD276 induces antitumor immunity as it activates both acquired and innate immunity leading to natural killer cell and CD8 T-cell dependent killing of tumor cells.

**PRESENTATION:**

**Purified:** Purified IgG buffered in PBS and 0.02% NaN<sub>3</sub>. (Purified via Protein G Chromatography). For maximum recovery of contents, spin down tube before use.

**LE:** Purified Ig buffered in PBS, no preservative, 0.2µm sterile filtered. (Purified from cell culture supernatant via Protein G Chromatography)

**Biotin, FITC, PE and APC:** Biotin/FITC/PE/APC conjugated IgG buffered in PBS, 0.02% NaN<sub>3</sub> and EIA grade BSA as a stabilizing protein to bring total protein concentration to 4-5 mg/ml.

**STORAGE/STABILITY:**

For all formats, store at 4°C. DO NOT FREEZE **PE and APC** conjugates. For long term storage (**Purified, LE, Biotin** and **FITC**), aliquot and freeze unused portion at -20°C in volumes appropriate for single usage. Avoid freeze/thaw cycles.

**APPLICATIONS:**

This antibody is suitable for use in Flow Cytometry and has also been shown to block CD276 when administered *in vivo*.

*Continued.....*

Visit our website for your local distributor.

**CEDARLANE**<sup>®</sup>

[www.cedarlanelabs.com](http://www.cedarlanelabs.com)

An ISO 9001:2015 and ISO 13485:2016  
registered company.

In CANADA: Toll Free: 1-800-268-5058

4410 Paletta Court, Burlington, ON L7L 5R2 ph: (289) 288-0001, fax: (289) 288-0020  
e-mail: [general@cedarlanelabs.com](mailto:general@cedarlanelabs.com)

In the USA: Toll Free: 1-800-721-1644

1210 Turrentine Street, Burlington, NC 27215 ph: (336) 513-5135, fax: (336) 513-5138  
e-mail: [service@cedarlanelabs.com](mailto:service@cedarlanelabs.com)

**SPECIFICATIONS:**

Clone: MJ18

Hybridoma Production:

Immunization: Immunogen: B7-H3-Ig, consisting of the extracellular domain (aa 1–242 of mouse B7-H3) linked to the Fc portion of mouse IgG2a.

Donor: Sprague Dawley Lymph node cells

Fusion Partner: P3U1 myeloma cell line.

Specificity: Mouse CD276 (B7-H3)

**REFERENCES:**

1. Nagashima, O. et al. (2008) B7-H3 contributes to the development of pathogenic Th2 cells in a murine model of asthma. *J Immunol.* 181 (6): 4062-71.
2. Fukushima, A. et al. (2007) B7-H3 regulates the development of experimental allergic conjunctivitis in mice. *Immunol Lett.* 113 (1): 52-7.

**FOR RESEARCH USE ONLY**  
® is a registered trademark of Cedarlane