



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

# TECHNICALLY *Speaking*

Place your order with CEDARLANE® or your local distributor.

*Please contact CEDARLANE® for lot specific information.*

## **FITC Anti-Mouse CD5 (Ly 1.2) Monoclonal Antibody**

**CL8912F  
CL8912F-3  
LOT: 8231**

### **DESCRIPTION:**

Cedarlane's anti-CD5 (Ly 1.2) mAb reacts with T cells from mouse strains expressing the Ly 1.2 phenotype, but does not react with lymphocytes from mouse strains expressing the Ly 1.1 phenotype.

### **PRESENTATION:**

100 µg (CL8912F) or 300 µg (CL8912F-3) FITC conjugated Ig buffered in PBS, 0.02% NaN<sub>3</sub> and EIA grade BSA as a stabilizing protein to bring total protein concentration to 4-5 mg/ml.

### **STORAGE/STABILITY:**

Store at 4°C. For long term storage, aliquot and freeze unused portion at -20°C in volumes appropriate for single usage. Avoid freeze/thaw cycles. Avoid prolonged exposure to light.

For more information or to place an order please contact...

**CEDARLANE®**  
**LABORATORIES LIMITED**



**toll free: 1-800-268-5058**  
*in North America*

**phone: (905) 878-8891 • fax: (905) 878-7800**

5516 - 8th Line, R.R.#2, Hornby, Ontario, CANADA L0P 1E0

or visit our website for a list of our international distributors including contact information

**website: [www.cedarlanelabs.com](http://www.cedarlanelabs.com) • e-mail: [info@cedarlanelabs.com](mailto:info@cedarlanelabs.com)**

**SPECIFICATIONS:**

Clone: CG16

**Hybridoma Production:**

Immunization: Immunogen: C3H.CE - Ly 1.2 : DS

Donor: C3H spleen

Fusion Partner: Myeloma SP2/0 - Ag 14 (M5).

Specificity: Mouse CD5 (Ly 1.2)

Ig Class: Mouse IgG<sub>2b</sub>

Format: FITC conjugated Ig buffered in PBS, 0.02% NaN<sub>3</sub> and EIA grade BSA as a stabilizing protein to bring total protein concentration to 4-5 mg/ml. (Purified from ascitic fluid via Protein G Chromatography)

Antibody Concentration: 0.1 mg/ml

**FLOW CYTOMETRY ANALYSIS:****Method:**

1. Prepare a cell suspension in media A. For cell preparations, deplete the red blood cell population with Lympholyte<sup>®</sup>-M cell separation medium (CL5030).
2. Wash 2 times.
3. Resuspend the cells to a concentration of  $2 \times 10^7$  cells/ml in media A. Add 50  $\mu$ l of this suspension to each tube (each tube will then contain  $1 \times 10^6$  cells, representing 1 test).
4. To each tube, add 0.1 - 0.2  $\mu$ g\* of **CL8912F** or **CL8912F-3** per  $10^6$  cells.
5. Vortex the tubes to ensure thorough mixing of antibody and cells.
6. Incubate the tubes for 30 minutes at 4°C.
7. Wash 2 times at 4°C.
8. Resuspend the cell pellet in 50  $\mu$ l ice cold media B.
9. Transfer to suitable tubes for flow cytometric analysis containing 15  $\mu$ l of propidium iodide at 0.5 mg/ml in PBS. This stains dead cells by intercalating in DNA.

Media:

- A. Phosphate buffered saline (pH 7.2) + 5% normal serum of host species + sodium azide (100 µl of 2M sodium azide in 100 mls).
- B. Phosphate buffered saline (pH 7.2) + 0.5% Bovine serum albumin + sodium azide (100 µl of 2M sodium azide in 100 mls).

Results:Tissue Distribution by Flow Cytometry Analysis:

Mouse Strain: BALB/c


Cell Concentration :  $1 \times 10^6$  cells per tests

Antibody Concentration Used:  $0.2 \mu\text{g}/10^6$  cells

Isotypic Control: FITC Mouse IgG<sub>2b</sub>

Cell SourcePercentage of cells stained above control:

Thymus	98.9%
Spleen	33.5%
Lymph Node	88.6%
Bone Marrow	3.3%



Cell Source: Thymus

Percentage of cells stained above control: 98.9 %

**N.B. Appropriate control samples should always be included in any labelling studies.**

**\* For optimal results in various applications, it is recommended that each investigator determine dilutions appropriate for individual use.**

Strain Distribution by Flow Cytometry Analysis:

Procedure: see page 2

Cell Concentration :  $1 \times 10^6$  cells per tests

Antibody Concentration Used:  $0.2 \mu\text{g}/10^6$  cells

Strains Tested: AKR, ATH, BALB/c, CBA/J, C3H/He

Positive: AKR, ATH, BALB/c

Negative: CBA/J, C3H/He

**FOR RESEARCH USE ONLY**

® is a Registered Trademark of Cedarlane Laboratories Limited.