



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

TECHNICALLY *Speaking*

Place your order with CEDARLANE® or your local distributor.

Please contact CEDARLANE® for lot specific information.

PE Anti-Mouse CD5 (Ly 1.2) Monoclonal Antibody

CL8912PE
CL8912PE-3
LOT: 8253

DESCRIPTION:

Cedarlane's anti-CD5 (Ly 1.2) monoclonal antibody reacts with T cells from mouse strains expressing the Ly 1.2 phenotype, but does not react with lymphocytes from mouse strains expressing the Ly 1.1 phenotype.

PRESENTATION:

50 µg (CL8912PE) or 300 µg (CL8912PE-3) PE conjugated Ig buffered in PBS, 0.02% NaN₃ and EIA grade BSA as a stabilizing protein to bring total protein concentration to 4-5 mg/ml.

STORAGE/STABILITY:

Store at 4°C. **DO NOT FREEZE.** Avoid prolonged exposure to light.

For more information or to place an order please contact...

CEDARLANE®
LABORATORIES LIMITED



toll free: 1-800-268-5058
in North America

phone: (905) 878-8891 • fax: (905) 878-7800

5516 - 8th Line, R.R.#2, Hornby, Ontario, CANADA L0P 1E0

or visit our website for a list of our international distributors including contact information

website: www.cedarlanelabs.com • e-mail: info@cedarlanelabs.com

SPECIFICATIONS:

Clone: CG16

Hybridoma Production:

Immunization: Immunogen: C3H.CE - Ly 1.2 : DS
Donor: C3H spleen
Fusion Partner: Myeloma SP2/0 - Ag 14 (M5).

Specificity: Mouse CD5 (Ly 1.2)

Ig Class: Mouse IgG_{2b}

Format: R-PE conjugated purified Ig buffered in PBS, 0.02% NaN₃ and EIA grade BSA as a stabilizing protein to bring total protein concentration to 4-5 mg/ml. (Purified from ascitic fluid via Protein G Chromatography)

Antibody Concentration: 0.1 mg/ml

FLOW CYTOMETRY ANALYSIS:**Method:**

1. Prepare a cell suspension in media A. For cell preparations, deplete the red blood cell population with Lympholyte[®]-M cell separation medium (CL5030).
2. Wash 2 times.
3. Resuspend the cells to a concentration of 2×10^7 cells/ml in media A. Add 50 μ l of this suspension to each tube (each tube will then contain 1×10^6 cells, representing 1 test).
4. To each tube, add 0.5 μ g* of **CL8912PE or CL8912PE-3** per 10^6 cells.
5. Vortex the tubes to ensure thorough mixing of antibody and cells.
6. Incubate the tubes for 30 minutes at 4°C.
(It is recommended that the tubes are protected from light, since most fluorochromes are light sensitive.)
7. Wash 2 times at 4°C.
8. Resuspend the cell pellet in 50 μ l ice cold media B.
9. Transfer to suitable tubes for flow cytometric analysis containing 15 μ l of propidium iodide at 0.5 mg/ml in PBS. This stains dead cells by intercalating in DNA.

Media:

- A. Phosphate buffered saline (pH 7.2) + 5% normal serum of host species + sodium azide (100 µl of 2M sodium azide in 100 mls).
- B. Phosphate buffered saline (pH 7.2) + 0.5% Bovine serum albumin + sodium azide (100 µl of 2M sodium azide in 100 mls).

Results:Tissue Distribution by Flow Cytometry Analysis:

Mouse Strain: BALB/c


Cell Concentration : 1×10^6 cells per tests

Antibody Concentration Used: $0.5 \mu\text{g}/10^6$ cells

Isotypic Control: PE Mouse IgG_{2b}

Cell SourcePercentage of cells stained above control:

Thymus	98.8%
Spleen	18.7%
Lymph Node	65.4%
Bone Marrow	11.3%



LFL2

Cell Source: Thymus

Percentage of cells stained above control: 98.8 %

N.B. Appropriate control samples should always be included in any labelling studies.

***For optimal results in various applications, it is recommended that each investigator determine dilutions appropriate for individual use.**

R-Phycoerythrin conjugates are produced under license and protected under Stanford University held patents 4,520,110; 4,542,104; 4,859,582; 5,055,556 (U.S.); 76695 (EPC); 548440 (Australia); 1,179,942 (Canada); and 1,594,827 (Japan).

Strain Distribution by Flow Cytometry Analysis:

Procedure: see page 2

Cell Concentration : 1×10^6 cells per tests

Antibody Concentration Used: $0.5 \mu\text{g}/10^6$ cells

Strains Tested: AKR, ATH, BALB/c, CBA/J, C3H/He

Positive: AKR, ATH, BALB/c

Negative: CBA/J, C3H/He

FOR RESEARCH USE ONLY

® is a Registered Trademark of Cedarlane Laboratories Limited.