



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

TECHNICALLY *Speaking*

Place your order with CEDARLANE® or your local distributor.

Please contact CEDARLANE® for lot specific information.

FITC Anti-Mouse CD62L Monoclonal Antibody

CL8918F
CL8918F-3
LOT: 1831

DESCRIPTION:

Cedarlane's anti-mouse CD62L (L-selectin Ly 22) monoclonal antibody reacts with a 90 kDa protein which is involved with the homing of lymphocytes to peripheral lymph nodes.

L-selectin is expressed on most T and B lymphocytes, neutrophils, monocytes, eosinophils.¹ Pre-incubation of lymphocytes with this antibody completely and specifically blocks binding of lymphocytes to high endothelial venules (HEV) in vitro^{2,3,6} and the migration of lymphocytes to lymph nodes in vivo.^{2,3} Polymorphonuclear cells preincubated with this antibody do not migrate to the inflammatory foci.³

Applications of CL8918F include flow cytometry^{3,4,5} and IHC.⁴

PRESENTATION:

100 µg (CL8918F) or 300 µg (CL8918F-3) FITC conjugated Ig buffered in PBS, 0.02% NaN₃ and EIA grade BSA as a stabilizing protein to bring total protein concentration to 4-5 mg/ml.

STORAGE/STABILITY:

Store at 4°C. For long term storage, aliquot and freeze unused portion at -20°C in volumes appropriate for single usage. Avoid freeze/thaw cycles. Avoid prolonged exposure to light.

For more information or to place an order please contact...

CEDARLANE®
LABORATORIES LIMITED



toll free: 1-800-268-5058
in North America

phone: (905) 878-8891 • fax: (905) 878-7800

5516 - 8th Line, R.R.#2, Hornby, Ontario, CANADA L0P 1E0

or visit our website for a list of our international distributors including contact information
website: www.cedarlanelabs.com • e-mail: info@cedarlanelabs.com

SPECIFICATIONS:

Clone: MEL-14

Hybridoma Production:

Immunization: Immunogen: Mouse B cell Lymphoma, 38C-14

Donor: Fischer Rat Spleen

Fusion Partner: P3 X 63Ag8.653

Specificity: Mouse CD62L (L-selectin Ly 22)

Ig Class: Rat IgG_{2a}

Format: FITC conjugated Ig buffered in PBS, 0.02% NaN₃ and EIA grade BSA as a stabilizing protein to bring total protein concentration to 4-5 mg/ml. (Purified from ascitic fluid via Protein G Chromatography)

Antibody Concentration: 0.1 mg/ml

FLOW CYTOMETRY ANALYSIS:

Method:

1. Prepare a cell suspension in media A. For cell preparations, deplete the red blood cell population with Lympholyte[®]-M cell separation medium (CL5030).
2. Wash 2 times.
3. Resuspend the cells to a concentration of 2×10^7 cells/ml in media A. Add 50 μ l of this suspension to each tube (each tube will then contain 1×10^6 cells, representing 1 test).
4. To each tube, add 0.2 - 0.5 μ g* of **CL8918F** or **CL8918F-3** per 10^6 cells.
5. Vortex the tubes to ensure thorough mixing of antibody and cells.
6. Incubate the tubes for 30 minutes at 4°C.
(It is recommended that the tubes are protected from light, since most fluorochemicals are light sensitive).
7. Wash 2 times at 4°C.
8. Resuspend the cell pellet in 50 μ l ice cold media B.
9. Transfer to suitable tubes for flow cytometric analysis containing 15 μ l of propidium iodide at 0.5 mg/ml in PBS. This stains dead cells by intercalating in DNA.

Media:

- A. Phosphate buffered saline (pH 7.2) + 5% normal serum of host species + sodium azide (100 µl of 2M sodium azide in 100 mls).
- B. Phosphate buffered saline (pH 7.2) + 0.5% Bovine serum albumin + sodium azide (100 µl of 2M sodium azide in 100 mls).

Results:Tissue Distribution by Flow Cytometry Analysis:

Mouse Strain: BALB/c


Cell Concentration : 1×10^6 cells per test

Antibody Concentration Used: $0.2 \mu\text{g}/10^6$ cells

Isotypic Control: FITC Rat IgG_{2a}

Cell SourcePercentage of cells stained above control:

Thymus	88.8%
Spleen	36.6%
Lymph Node	74.4%



Cell Source: Lymph Node

Percentage of cells stained above control: 74.4%

N.B. Appropriate control samples should always be included in any labelling studies.

*** For optimal results in various applications, it is recommended that each investigator determine dilutions appropriate for individual use.**

Strain Distribution by Flow Cytometry Analysis:

Procedure: see page 2

Cell Concentration: 1×10^6 cells per test

Antibody Concentration Used: $0.2 \mu\text{g}/10^6$ cells

Strains Tested: BALB/c, CBA/J, C3H/He, C57BL/6, AKR

Positive: BALB/c, CBA/J, C3H/He, C57BL/6, AKR

Negative: none

REFERENCES:

- 1) Fink, P., W. Gallatin, R. Reichert, *et al.* 1985. Homing receptor-bearing thymocytes, an immunocomponent cortical subpopulation. *Nature* 313: 233-235
- 2) Gallatin, W.M., I.L. Weissman., E.C. Butcher 1983. A cell surface molecule involved in organ specific homing of lymphocytes. *Nature* 304:30-34
- 3) Lewinsohn, D.M., R.F. Bargatze, E.C. Butcher 1987. Leukocyte-endothelial cell recognition: evidence of a common molecular mechanism shared by neutrophils, lymphocytes and other leukocytes. *J.Immunology* 138:4313-4321
- 4) Reichert, R., M. Gallatin, E. Butcher, *et al.*. 1984. A homing receptor-bearing cortical thymocyte subset: Implications for thymus cell migration and the nature of cortisone-resistant thymocytes. *Cell* 38: 89-99
- 5) Siegelman, M., I.C. Cheng, I.L. Weissman, *et al.* 1990. The mouse lymph node homing receptor is identical with the lymphocyte cell surface receptor Ly-22: Role of the EGF domain in endothelial binding. *Cell* 61: 611-622
- 6) Jalkanen, S., R.F. Bargatze, J. Toyos, *et al.* 1987. Lymphocyte Recognition of High Endothelium: Antibodies to Distinct Epitopes of an 85-95-kD Glycoprotein Antigen Differentially Inhibit Lymphocyte Binding to Lymph Node, Mucosal, or Synovial Endothelial Cell. *J. of Cell Biol.* 105: 983-990

FOR RESEARCH USE ONLY

® is a Registered Trademark of Cedarlane Laboratories Limited.