



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Technically
Speaking

CEDARLANE[®]
www.cedarlanelabs.com



Conveniently Delivering You Today's Innovations
for the Science of Tomorrow™

**Purified Anti-Mouse CD11b (MAC-1)
Monoclonal Antibody**

Catalogue#	Format	Size	Concentration	Isotype Control
CL8941A	Ascites	0.5 ml	NA	CLCR2B00
CL8941AP	Purified	250 µg	0.1 mg/ml	CLCR2B00
CL8941LE	Low Endotoxin	500 µg	1.0 mg/ml	CLCR2B00
CL8941B/-3	Biotin	100 µg/300 µg	0.1 mg/ml	CLCR2B15
CL8941F/-3	FITC	100 µg/300 µg	0.1 mg/ml	CLCR2B01
CL8941NA	No Azide	1.0 mg	1.0 mg/ml	CLCR2B00
CL8941PE/-3	PE	50 µg/300 µg	0.1 mg/ml	CLCR2B04
CL8941TC	PE-Cy5	100 µg	1.0 mg/ml	CLCR2B06
CL8941APC	APC	100 µg	0.1 mg/ml	CLCR2B05
CL8941G/-2	10 nm Gold	0.5 ml/1.0 ml	OD3	N/A

Ig Class: Rat IgG_{2b}

DESCRIPTION:

Cedarlane's anti-mouse CD11b (Mac-1; Ly 40) monoclonal antibody detects the 170 kDa α subunit of Mac-1 which mediates adhesion to ICAM-1 (CD54) and C3bi. Mac-1 is expressed on granulocytes, macrophages, natural killer cells, and B-1 cells in the peritoneal and pleural cavities. Mac-1 is up-regulated on neutrophils after activation. This particular clone blocks cell adherence and C3bi binding but does not block cell mediated lysis (1,2,3,4,5,6). Applications include flow cytometry, *in vitro* and *in vivo* blocking, immunohistochemistry for both paraffin embedded and acetone-fixed frozen sections 1-20 µg/ml (7). Also useful for immunoprecipitation and western blotting.

PRESENTATION:

Purified: Purified IgG buffered in PBS and 0.02% NaN₃. (Purified from ascitic fluid via Protein G Chromatography). For maximum recovery of contents, spin down tube before use.

Biotin, FITC, PE and PE-Cy5: Biotin/FITC/PE/PE-Cy5 conjugated IgG buffered in PBS, 0.02% NaN₃ and EIA grade BSA as a stabilizing protein to bring total protein concentration to 4-5 mg/mL.

No Azide: Purified Ig buffered in PBS, no preservative, 0.2 µm sterile filtered.

STORAGE/STABILITY:

For all formats, store at 4°C. DO NOT FREEZE PE, PE-Cy5, APC and Gold conjugates. For long term storage (**Purified, Biotin, FITC, No Azide**), aliquot and freeze unused portion at -20°C in volumes appropriate for single usage. Avoid freeze/thaw cycles.

Visit our website for your local distributor.

CEDARLANE[®]



www.cedarlanelabs.com

An ISO 9001:2000 and ISO 13485:2003
registered company.

In CANADA: Toll Free: 1-800-268-5058

4410 Paletta Court, Burlington, ON L7L 5R2 ph: (289) 288-0001, fax: (289) 288-0020
e-mail: general@cedarlanelabs.com

In the USA: Toll Free: 1-800-721-1644

1210 Turrentine Street, Burlington, NC 27215 ph: (336) 513-5135, fax: (336) 513-5138
e-mail: service@cedarlanelabs.com

SPECIFICATIONS:

Clone: M1/70.15

Hybridoma Production:

Immunization: Immunogen: C57BL/6 spleen cell enriched for T lymphocytes
Donor: DA rat spleen

Fusion Partner: NS-1

Specificity: Mouse CD11b (MAC-1, Ly-40)

TEST RESULTS:

Tissue Distribution by Flow Cytometry Analysis:

Mouse Strain: C₃H/HE

Cell Concentration: 1x10⁶ cells per test

Antibody Concentration Used: 0.2 µg/10⁶ cells

Cell Source

Peritoneal Macrophages

Bone Marrow Macrophages



Thymus

Percentage of cells stained above control:

90.4%

68.3%

7.5%



C3H/HE mouse peritoneal macrophages (left) and bone marrow (right) were stained with anti-CD11b (clone: M1/70.15) (filled histogram) or rat IgG2b isotype control (open histogram).

N.B. Appropriate control samples should always be included in any labeling studies.

*** For optimal results in various applications, it is recommended that each investigator determine dilutions appropriate for individual use.**

REFERENCES:

1. Sanchez-Madrid, F., P. Simon, S. Thompson, et al. 1983. Mapping of antigenic and functional epitopes on the α and β subunits of two related mouse glycoproteins involved in cell interactions, LFA-1 and Mac-1. *J. Exp. Med.* 158:586-602.
2. Springer, T., G. Galfre, D.S. Secher, et al. 1978 Monoclonal xenogeneic antibodies to murine cell surface antigens: identification of novel leukocyte differentiation antigens. *Eur. J. Immunol.* 8: 539 - 551.
3. Springer, T., G. Galfre, D.S. Secher, et al. 1979. Mac-1: a macrophage differentiation antigen identified by monoclonal antibody. *Eur. J. Immunol.* 9:301 - 306.
4. Vignali, D.A., et al. 1990. Antibody-Dependent killing of *Schistosoma mansoni* Schistosomula in vitro by Starch-elicited murine macrophages. *J. Immunol.* 144: 4030 -40378.
5. Kantor, A., et al. 1992. Differential development of progenitor activity for three B-cell lineages. *Proc. Natl. Acad. Sci. U.S.A.* 89: 3320 - 3324.
6. Jutial, M.A., et al. 1988. Ly-6C is a monocyte/macrophage and endothelial cell differentiation antigen regulated by interferon gamma. *J. Immunol.* 18: 1819 - 1826.
7. Whiteland, J.L. et al. (1995). Immunohistochemical detection of T cell subsets and other leukocytes in paraffin embedded rat and mouse tissues with monoclonal antibodies. *J. Histochem. Cytochem.* Volume 43, pp 313-320.