

# Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



# Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

# Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

# SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in







Conveniently Delivering You Today's Innovations for the Science of Tomorrow™

# **Anti-Mouse CD44 Monoclonal Antibody**

Catalogue#	Format	Size	Concentration
CL8946AP	Purified	200 μg	1.0 mg/ml
CL8946B	Biotin	100 μg	0.1  mg/ml
CL8946B-3	Biotin	300 μg	0.1 mg/ml
CL8946F	FITC	100 μg	0.1 mg/ml
CL8946F-3	FITC	300 μg	0.1 mg/ml
CL8946PE	PE	50 μg	0.1 mg/ml
CL8946PE-3	PE	300 μg	0.1 mg/ml

Isotype: Rat IgG<sub>2b</sub>

#### **DESCRIPTION:**

Cedarlane's anti-mouse CD44 monoclonal antibody reacts with all isoforms of CD44 (Pgp-1, Ly-24) glycoprotein. By flow cytometry, the main cellular reactivities are B cells, monocytes, macrophages and variable subsets of thymocytes and peripheral T cells.

This antibody is suitable for use in flow cytometry.

CL8946AP is reported to work well in immunohistochemistry (both paraffin (2.5-10ug/ml) and frozen sections). CL8946PE and CL8946PE-3 are reported to work in immunoprecipitation, ELISA, IHC of frozen sections, and complement depletion<sup>1,2,3</sup>.

### PRESENTATION:

**Purified:** 22 µg purified Ig buffered in PBS and 0.02% NaN<sub>3</sub>

Biotin and FITC: 100 µg or 300 µg Biotin/FITC conjugated Ig buffered in PBS, 0.02% NaN3 and EIA grade BSA as a stabilizing protein to bring total protein concentration to 4-5 mg/ml.

PE: 50 μg or 300 μg R-PE conjugated Ig buffered in PBS, 0.2% NaN3 and EIA grade BSA as a stabilizing protein to bring total protein concentration to 4-5 mg/ml.

## STORAGE/STABILITY:

Store at 4°C. Avoid prolonged exposure to light. DO NOT FREEZE PE conjugates. For long term storage (**PURIFIED, BIOTIN, FITC**), aliquot and freeze unused portion at -20°C in volumes appropriate for single usage. Avoid freeze/thaw cycles. If the reagent is being diluted, it is recommended that only the quantity to be used within one week be diluted. Check label for expiry date.

Visit our website for your local distributor.



In CANADA: Toll Free: 1-800-268-5058

4410 Paletta Court, Burlington, ON L7L 5R2 ph: (289) 288-0001, fax: (289) 288-0020 e-mail: general@cedarlanelabs.com

In the USA: Toll Free: 1-800-721-1644

1210 Turrentine Street, Burlington, NC 27215 ph: (336) 513-5135, fax: (336) 513-5138 e-mail: service@cedarlanelabs.com

### **SPECIFICATIONS:**

Clone: IM7.8.1

Specificity: Mouse CD44

### **TEST RESULTS:**

#### Tissue Distribution by Flow Cytometry Analysis:


Mouse Strain: BALB/c

Cell Concentration :  $1x10^6$  cells per test Antibody Concentration Used:  $0.5 \mu g/10^6$  cells Isotypic Control: FITC Rat  $IgG_{2b}$  (CLCR2B01)

<u>Cell Source:</u> <u>Percentage of cells stained above control:</u>

 Spleen
 76.5%

 Thymus
 20.9%



#### N.B. Appropriate control samples should always be included in any labeling studies.

\* For optimal results in various applications, it is recommended that each investigator determine dilutions appropriate for individual use.

#### **REFERENCES:**

- 1. Lesley, J. and Trowbridge, I.S. 1982 Genetic characterization of a polymorphic murine cell-surface glycprotein. Immunogenetics 15:313-320.
- 2. Lesley, J., Hyman, R. and Kincade, P.W. 1993 CD44 and its interaction with extracellular matrix. Advances in Immunol. 54:271-335.
- 3. Katoh, S., J.B. McCarthy, and P.W. Kincade. 1994. Characterization of soluble CD44 in the circulation of mice. Levels are affected by immune activity and tumor growth. J.Immunol. 153:3440-3449.

® is a Registered Trademark of Cedarlane Laboratories Limited.