

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in





Conveniently Delivering You Today's Innovations for the Science of Tomorrow™

Anti-Mouse/Rat CD81 **Monoclonal Antibody**

Catalogue#	Format	Size	Concentration	Isotype Control
CL8981AP	Purified	250µg	1.0 mg/ml	CLCHM00
CL8981NA	No Azide	1.0mg	1.0 mg/ml	CLCHM00
CL8981B/-3	Biotin	100μg/300μg	0.1 mg/ml	CLCHM15
CL8981F/-3/-5	FITC	100μg/300μg/500 μg	0.1 mg/ml	CLCHM01
CL8981PE/-3/-4	PE	50μg/300μg /200μg	0.1 mg/ml	CLCHM04
CL8981AF4	Alexa Fluor®488	100 μg	0.1 mg/ml	N/A

Alexa Fluor® is a registered trademark of Life Technologies Corporation.

Isotype: Armenian Hamster IgG

DESCRIPTION:

Cedarlane's CL8981 reacts with the extracellular loops of murine CD81 (TAPA-1) molecule. As a member of the tetraspanin superfamily of cell-surface proteins, CD81 has been linked to the control of cell proliferation, adhesion and motility. CD81 is expressed in higher levels on resting murine B cells than on resting T cells and is functionally active on B cells as it induces homotypic adhesion of B lymphocytes. Unlike human CD81, which is expressed equally on all thymocytes, murine CD81 is upregulated on CD4+CD8+ thymocytes, then down-regulated again on mature single-positive thymocytes. Murine dendritic cells, splenic macrophages and non-killer cells all express very high levels of CD81. CD81 has also been involved in the induction of IL-4 secretion from T cells during Th2 immune responses. It has been reported that CD81 expression can also be induced in mature T cells upon activation. This anti-CD81 mAb has been shown to decrease the proliferation of LPS stimulated CD81^{+/+} B cells to levels similar to that of CD81^{-/-} B cells.

PRESENTATION:

Purified: Purified IgG buffered in PBS and 0.02% NaN3. (Purified from ascitic fluid via Protein G Chromatography). For maximum recovery of contents, spin down tube before use.

Biotin, FITC, PE and AF488: Biotin/FITC/PE/AF488 conjugated IgG buffered in PBS, 0.02% NaN3 and EIA grade BSA as a stabilizing protein to bring total protein concentration to 4-5 mg/ml.

No Azide: Purified Ig buffered in PBS,no preservative,0.2µm sterile filtered.

STORAGE/STABILITY:

For all other formats, store at 4°C. DO NOT FREEZ PE and AF488 conjugates. For long term storage (Purified, Biotin, NA and FITC), aliquot and freeze unused portion at -20°C in volumes appropriate for single usage. Avoid freeze/thaw cycles.

Visit our website for your local distributor.



In CANADA: Toll Free: 1-800-268-5058

4410 Paletta Court, Burlington, ON L7L 5R2 ph: (289) 288-0001, fax: (289) 288-0020 e-mail: general@cedarlanelabs.com

In the USA: Toll Free: 1-800-721-1644

1210 Turrentine Street, Burlington, NC 27215 ph: (336) 513-5135, fax: (336) 513-5138 e-mail: service@cedarlanelabs.com

SPECIFICATIONS:

Clone: Eat2

Immunogen: CD81+ mouse B lymphoma 38C13


Specificity: Mouse CD81 (TAPA-1)

TEST RESULTS:

<u>Tissue distribution by Flow Cytometry Analysis:</u>

Mouse Strain: C57BL/6

Cell Concentration: 1×10^6 cells per test Antibody Concentration: $1.0 \mu g/10^6$ cells



Cell Source: Mouse Splenic B Cells**
Percentage of cells stained above control: 92.7%

N.B. Appropriate control samples should always be included in any labeling studies.

* For optimal results in various applications, it is recommended that each investigator determine dilutions appropriate for individual use.

REFERENCES:

- 1. Maecker, H. T., et al., 2000. Differential expression of murine CD81 highlighted by new anti-mouse CD81 monoclonal antibodies. *Hybridoma* 19:15-22.
- 2. Maecker, H. T., Todd, S. C., and Levy, S, 1997. The tetraspanin superfamily: Molecular facilitators. *Faseb J* 11:428-442.
- 3. Miyazaki, T., Muller, U. and Campbell, K. S., 1997. Normal development but differentially altered proliferative responses of lymphocytes in mice lacking CD81. *Embo J* 16:4217-4225.

^{**} B cells isolated with CL131 - Cedarlane's Mouse B Cell Recovery Kit