



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

# TECHNICALLY *Speaking*

Place your order with CEDARLANE® or your local distributor.

*Please contact CEDARLANE® for lot specific information.*

## **Purified Anti-Mouse CD86 (B7-2) Monoclonal Antibody**

**CL8986AP**

**LOT:**

### **DESCRIPTION:**

Cedarlane's anti-mouse B7-2 monoclonal antibody reacts with a co-stimulatory molecule expressed on antigen-presenting cells, dendritic cells and activated B cells within 24 hours of stimulation by LPS, anti-CD40, anti-surface Ig, or specific antigen. B7-2, a ligand of CD28 and CTLA-4, is one of the accessory molecules that plays an important role in T cell-B cell co-stimulatory interactions.

This antibody is suitable for use in flow cytometry.

### **PRESENTATION:**

200 µg purified Ig buffered in PBS and 0.1% NaN<sub>3</sub>

### **STORAGE/STABILITY:**

Store at 4°C. For long term storage, aliquot and freeze unused portion at -20°C in volumes appropriate for single usage. Avoid freeze/thaw cycles. If the reagent is being diluted, it is recommended that only the quantity to be used within one week be diluted. Check label for expiry date.

For more information or to place an order please contact...

**CEDARLANE®**  
**LABORATORIES LIMITED**



*toll free: 1-800-268-5058  
in North America*

phone: (905) 878-8891 • fax: (905) 878-7800

5516 - 8th Line, R.R.#2, Hornby, Ontario, CANADA L0P 1E0

or visit our website for a list of our international distributors including contact information

website: [www.cedarlanelabs.com](http://www.cedarlanelabs.com) • e-mail: [info@cedarlanelabs.com](mailto:info@cedarlanelabs.com)

**SPECIFICATIONS:**

Clone: RMMP-2

Specificity: Mouse CD86 (B7-2)

Ig Class: Rat IgG<sub>2a</sub>

Format: Purified Ig buffered in PBS, 0.1% NaN<sub>3</sub> and EIA grade BSA to bring total protein concentration to 4-5 mg/ml.

Antibody Concentration: 0.2 mg/ml

**FLOW CYTOMETRY ANALYSIS:**

Method:

1. Prepare a cell suspension in media A. For cell preparations, deplete the red blood cell population with Lympholyte<sup>®</sup>-M cell separation medium (CL5030).
2. Wash 2 times.
3. Resuspend the cells to a concentration of  $2 \times 10^7$  cells/ml in media A. Add 50  $\mu$ l of this suspension to each tube (each tube will then contain  $1 \times 10^6$  cells, representing 1 test).
4. To each tube, add  $\sim 1.0 \mu\text{g}^*$  of **CL8986AP**.
5. Vortex the tubes to ensure thorough mixing of antibody and cells.
6. Incubate the tubes for 30 minutes at 4°C.
7. Wash 2 times at 4°C.
8. Add 100  $\mu$ l of secondary antibody **CLCC40001** (FITC Goat anti-rat IgG (H+L)) at 1:500 dilution.
9. Incubate the tubes at 4°C for 30-60 minutes.  
(It is recommended that the tubes are protected from light since most fluorochromes are light sensitive).
10. Wash 2 times at 4°C in media B.
11. Resuspend the cell pellet in 50  $\mu$ l ice cold media B.
12. Transfer to suitable tubes for flow cytometric analysis containing 15  $\mu$ l of propidium iodide at 0.5 mg/ml in PBS. This stains dead cells by intercalating in DNA.

Media:

- A. Phosphate buffered saline (pH 7.2) + 5% normal serum of host species + sodium azide (100 µl of 2M sodium azide in 100 mls).
- B. Phosphate buffered saline (pH 7.2) + 0.5% Bovine serum albumin + sodium azide (100 µl of 2M sodium azide in 100 mls).

**N.B. Appropriate control samples should always be included in any labelling studies.**

**\* For optimal results in various applications, it is recommended that each investigator determine dilutions appropriate for individual use.**

**FOR RESEARCH USE ONLY**

® is a Registered Trademark of Cedarlane Laboratories Limited.