



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Technically  
Speaking

**CEDARLANE**<sup>®</sup>  
[www.cedarlanelabs.com](http://www.cedarlanelabs.com)



Conveniently Delivering You Today's Innovations  
for the Science of Tomorrow™

**Anti-Mouse Neutrophils (Ly-6B.2)  
Monoclonal Antibody**

Catalogue#	Format	Size	Concentration	Isotype Control
<b>CL8993AP</b>	Purified	100µg	1.0 mg/ml	CLCR2A00
<b>CL8993B/-3</b>	Biotin	100µg/300µg	0.1 mg/ml	CLCR2A15
<b>CL8993F/-3</b>	FITC	100µg/300µg	0.1 mg/ml	CLCR2A01
<b>CL8993PE/-3</b>	PE	50µg/300µg	0.1 mg/ml	CLCR2A04

Isotype: Rat IgG<sub>2a</sub>

**DESCRIPTION:**

Cedarlane's anti-mouse neutrophils monoclonal antibody is specific for mouse neutrophils.

This antibody is suitable for use in flow cytometry. This clone has also been reported to work in immunohistochemistry (both frozen and paraffin sections)<sup>2</sup>. Strains reported to be positive for the 7/4 clone are: AKR, C57BL/6, C57BL/10, C58, DBA/2, MF1, NZB, NZW, SJL, Swiss (PO) and 129J. Strains reported to be negative/weak for the 7/4 clone are: A2G, A/Sn, ASW, BALB/c, C3H/HEH and CBA.T6T6.

**PRESENTATION:**

**Purified:** Purified IgG buffered in PBS and 0.09% NaN<sub>3</sub>. (Purified from tissue culture supernatant via Protein G Chromatography). For maximum recovery of contents, spin down tube before use.

**Biotin, FITC and PE:** Biotin/FITC/PE conjugated IgG buffered in PBS, 0.09% NaN<sub>3</sub> and EIA grade BSA as a stabilizing protein to bring total protein concentration to 4-5 mg/ml.

**STORAGE/STABILITY:**

Store all formats at 4°C. DO NOT FREEZE

**SPECIFICATIONS:**

Clone: 7/4

Immunogen: Cultured bone marrow cells

Specificity: Mouse Neutrophils

*Continued Overleaf.....*

Visit our website for your local distributor.

**CEDARLANE**<sup>®</sup>



[www.cedarlanelabs.com](http://www.cedarlanelabs.com)

An ISO 9001:2000 and ISO 13485:2003  
registered company.

**In CANADA: Toll Free: 1-800-268-5058**

4410 Paletta Court, Burlington, ON L7L 5R2 ph: (289) 288-0001, fax: (289) 288-0020  
e-mail: [general@cedarlanelabs.com](mailto:general@cedarlanelabs.com)

**In the USA: Toll Free: 1-800-721-1644**

1210 Turrentine Street, Burlington, NC 27215 ph: (336) 513-5135, fax: (336) 513-5138  
e-mail: [service@cedarlanelabs.com](mailto:service@cedarlanelabs.com)


## **TEST RESULTS:**

### Tissue Distribution by Flow Cytometry Analysis:

Mouse Strain: C57BL/6

Cell Concentration:  $1 \times 10^6$  cells per test

Antibody Concentration Used: 0.2 - 1.0  $\mu\text{g}/10^6$  cells



C57BL/6 mouse peripheral blood granulocytes were stained with anti-Neutrophils (Ly-6B.2) (clone: 7/4) (filled histogram) or rat IgG2a isotype control (open histogram).

**N.B. Appropriate control samples should always be included in any labeling studies.**

**\* For optimal results in various applications, it is recommended that each investigator determine dilutions appropriate for individual use.**

## **REFERENCES:**

1. Hirsch, S. and Gordon, S. (1983). Polymorphic expression of a neutrophil differentiation antigen revealed by a monoclonal antibody 7/4. *Immunogenetics* 18: 229-239.
2. Gordon, S. et al (1992). Antigen markers of macrophage differentiation in murine tissues. *Curr. Opin. Microbiol. Immunol.* 181: 1-37.

**Laboratory Reagent For Research Use Only**

® is a Registered Trademark of Cedarlane