



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

# TECHNICALLY *Speaking*

Place your order with CEDARLANE® or your local distributor.

*Please contact CEDARLANE® for lot specific information.*

## **FITC Anti-Mouse CD45RC Monoclonal Antibody**

**CL8997F**

**CL8997F-3**

**LOT: 9731**

### **DESCRIPTION:**

CD45 (L-CA) is a transmembrane phosphotyrosine phosphatase expressed on leukocytes. Cedarlane's anti-mouse CD45RC is against the exon C-dependent RC isoform and reacts strongly with B cells, and less intensely with most CD8+ T cells. It does not recognize CD4+ T cells. Also, myeloid cells do not express the RC isoform.

Applications for this clone include immunohistochemistry (acetone-fixed frozen sections), flow cytometry and immunoprecipitation

### **PRESENTATION:**

100 µg (CL8997F) or 300 µg (CL8997F-3) FITC conjugated Ig buffered in PBS, 0.02% NaN<sub>3</sub> and EIA grade BSA as a stabilizing protein to bring total protein concentration to 4-5 mg/ml.

### **STORAGE/STABILITY:**

Store at 4°C. For long term storage, aliquot and freeze unused portion at -20°C in volumes appropriate for single usage. Avoid freeze/thaw cycles.

For more information or to place an order please contact...

**CEDARLANE®**  
**LABORATORIES LIMITED**



*toll free: 1-800-268-5058  
in North America*

phone: (905) 878-8891 • fax: (905) 878-7800

5516 - 8th Line, R.R.#2, Hornby, Ontario, CANADA L0P 1E0

or visit our website for a list of our international distributors including contact information

website: [www.cedarlanelabs.com](http://www.cedarlanelabs.com) • e-mail: [info@cedarlanelabs.com](mailto:info@cedarlanelabs.com)

## **SPECIFICATIONS:**

Clone: IBL-8

### Hybridoma Production:

Immunization: Immunogen: Mouse Spleen Cells  
Donor: Wistar spleen

Fusion Partner: Sp-2/0 Ag14

Specificity: Mouse CD45RC

Ig Class: : Rat IgG<sub>1</sub>/k

Format: FITC conjugated Ig buffered in PBS, 0.02% NaN<sub>3</sub> and EIA grade BSA as a stabilizing protein to bring total protein concentration to 4-5 mg/ml.

Antibody Concentration: 0.1 mg/ml

## **FLOW CYTOMETRY ANALYSIS:**

### Method:

1. Prepare a cell suspension in media A. For cell preparations, deplete the red blood cell population with Lympholyte<sup>®</sup>-M cell separation medium (CL5030).
2. Wash 2 times.
3. Resuspend the cells to a concentration of  $2 \times 10^7$  cells/ml in media A. Add 50  $\mu$ l of this suspension to each tube (each tube will then contain  $1 \times 10^6$  cells, representing 1 test).
4. To each tube, add 0.5  $\mu$ g of **CL8997F or CL8997F-3** per  $10^6$  cells.
5. Vortex the tubes to ensure thorough mixing of antibody and cells.
6. Incubate the tubes for 30 minutes at 4°C.  
(It is recommended that the tubes are protected from light, since most fluorochromes are light sensitive.)
7. Wash 2 times at 4°C.
8. Resuspend the cell pellet in 50  $\mu$ l ice cold media B.
9. Transfer to suitable tubes for flow cytometric analysis containing 15  $\mu$ l of propidium iodide at 0.5 mg/ml in PBS. This stains dead cells by intercalating in DNA.

### Media:

- A. Phosphate buffered saline (pH 7.2) + 5% normal serum of host species + sodium azide (100  $\mu$ l of 2M sodium azide in 100 mls).
- B. Phosphate buffered saline (pH 7.2) + 0.5% Bovine serum albumin + sodium azide (100  $\mu$ l of 2M sodium azide in 100 mls).

Results:

Tissue Distribution by Flow Cytometry Analysis:

Mouse Strain: BALB/c

Cell Concentration :  $1 \times 10^6$  cells per test

Antibody Concentration Used:  $0.5 \mu\text{g}/10^6$  cells


Isotypic Control: FITC Rat IgG<sub>1</sub> (CLCR101)

Cell Source

Percentage of cells stained above control:

Lymph Nodes

44.3%



Percentage of cells stained above control: 44.3 %

**N.B. Appropriate control samples should always be included in any labeling studies.**

**\* For optimal results in various applications, it is recommended that each investigator determine dilutions appropriate for individual use.**

**REFERENCES:**

1. Czompoly T, Labadi A, Balazs M, Nemeth P, and P Balogh. Use of cyclic peptide phage display library for the identification of a CD45RC epitope expressed on murine B cells and their precursors. *Biochemical and Biophysical Research Communications*. 2003 307: 791-796.

**FOR RESEARCH USE ONLY**

® is a Registered Trademark of Cedarlane Laboratories Limited.