



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

# TECHNICALLY *Speaking*

Place your order with CEDARLANE® or your local distributor.

*Please contact CEDARLANE® for lot specific information.*

## **FITC Anti-Mouse H-2K<sup>b</sup>D<sup>b</sup> Monoclonal Antibody**

**CL9007F**

**LOT:**

### **DESCRIPTION:**

Cedarlane's anti-mouse H-2K<sup>b</sup>D<sup>b</sup> monoclonal antibody is specific for cells expressing the H-2K antigen coded for by the b haplotype and for cells expressing the H-2D antigen coded for by the b haplotype.

### **PRESENTATION:**

100 µg FITC-labelled Ig buffered in PBS +0.02% NaN<sub>3</sub>. BSA was added to bring total protein concentration to 4-5 mg/ml.

### **STORAGE/STABILITY:**

Stable at 4°C. For long term storage, aliquot and freeze unused portions at -20°C in volumes appropriate for single usage. Avoid repeated freeze/thaw cycles. Avoid prolonged exposure to light.

For more information or to place an order please contact...

**CEDARLANE®**  
**LABORATORIES LIMITED**



*toll free: 1-800-268-5058  
in North America*

phone: (905) 878-8891 • fax: (905) 878-7800

5516 - 8th Line, R.R.#2, Hornby, Ontario, CANADA L0P 1E0

or visit our website for a list of our international distributors including contact information

website: [www.cedarlanelabs.com](http://www.cedarlanelabs.com) • e-mail: [info@cedarlanelabs.com](mailto:info@cedarlanelabs.com)

**SPECIFICATIONS:**

Clone: 5041.16.1

**Hybridoma Production:**

Immunization: Recipient: CBA/J  
Immunocyte Donor: bm12:thymus, lymph node, spleen

Fusion Partner: NS-1 mouse myeloma cell line

Specificity: Mouse H-2K<sup>b</sup>D<sup>b</sup>

Ig Class: Mouse IgG<sub>2a</sub>

Presentation: FITC-labelled Ig buffered in PBS with the addition of 0.02% NaN<sub>3</sub> and EIA grade BSA as a stabilizing protein to bring total protein concentration to 4-5 mg/ml. (FITC-labelled from ascitic fluid via Protein G Chromatography)

Antibody Concentration: 0.1 mg / ml

**FLOW CYTOMETRY ANALYSIS:**

1. Prepare cell suspension in Media A. For cell preparations, deplete the red blood cell population with Lympholyte<sup>®</sup>-M cell separation media (CL5030).
2. Wash 2 times.
3. Resuspend the cells to a concentration of  $2 \times 10^7$  cells/ml in media A. Add 50  $\mu$ l of this suspension to each tube (each tube will then contain  $1 \times 10^6$  cells, representing 1 test).
4. To each tube add 0.1  $\mu$ g of **CL9007F** per  $10^6$  cells\*.
5. Vortex the tubes to ensure thorough mixing of antibody and cells.
6. Incubate the tubes for 30 minutes at 4°C.
7. Wash 2 times at 4°C.
8. Resuspend the cell pellet in 50  $\mu$ l ice cold Media B.
9. Transfer to suitable tubes for flow cytometric analysis containing 15  $\mu$ l of propidium iodide at 0.5 mg/ml in phosphate buffered saline. (This stains dead cells by intercalating DNA.)

**MEDIA:**

- A. Phosphate buffered saline (pH 7.2) + 5% normal serum of host species + sodium azide (100  $\mu$ l of 2 M sodium azide in 100 mls).
- B. Phosphate buffered saline (pH 7.2) + 0.5% bovine serum albumin + sodium azide (100  $\mu$ l of 2 M sodium azide in 100 mls).


**FLOW CYTOMETRIC ANALYSIS:**

Donor: C57BL/6

Cell Concentration:  $1 \times 10^6$  cells

Antibody Concentration: 0.1  $\mu$ g/ $10^6$  cells

Isotypic Control: FITC Mouse IgG<sub>2a</sub>  $\kappa$

**LFL1**

Cell Source: Spleen

Percentage of Cells Stained Above Control: 99.9%

**N.B.** Appropriate control samples should always be included in any labelling studies.

**\* For optimal results in various applications, it is recommended that each investigator determine dilutions appropriate for individual use.**

**STRAIN DISTRIBUTION:**

Procedure: As above

Antibody Concentration: 0.1µg/10<sup>6</sup> cells

Strains Tested:

<u>Strain</u>	<u>Haplotype</u>	<u>+/-</u>
BALB/c	H-2 <sup>d</sup>	-
C3H/He	H-2 <sup>k</sup>	-
CBA/J	H-2 <sup>k</sup>	-
C57BL/6	H-2 <sup>b</sup>	+
B6Lyt 2.1 3.1	H-2 <sup>b</sup>	+

**FOR RESEARCH USE ONLY**

(Not to be administered to humans or animals nor used for any drug purpose)

® is a registered Trademark of Cedarlane Laboratories Ltd.