



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Technically  
Speaking

**CEDARLANE**<sup>®</sup>  
[www.cedarlanelabs.com](http://www.cedarlanelabs.com)



Conveniently Delivering You Today's Innovations  
for the Science of Tomorrow™

**Anti-Mouse H-2K<sup>d</sup>D<sup>d</sup>  
Monoclonal Antibody**

Catalogue#	Format	Size	Concentration	Isotype Control
CL9023A	Ascites	0.5 ml	lyophilized	CLCMG2A00
CL9023AP	Purified	250µg	1.0 mg/ml	CLCMG2A00
CL9023B	Biotin	100µg	0.1 mg/ml	CLCMG2A15
CL9023F	FITC	100µg	0.1 mg/ml	CLCMG2A01
CL9023PE	PE	50µg	0.1 mg/ml	CLCMG2A04

Isotype: Mouse IgG<sub>2a</sub>

**DESCRIPTION:**

Cedarlane anti H-2K<sup>d</sup>D<sup>d</sup> mAb reacts with both H-2K<sup>d</sup> and H-2D<sup>d</sup> products. The antibody also cross-reacts with K<sup>bsr</sup>pq. This K-D cross reaction indicates the presence of shared specificities between the two separate H-2 regions.

**PRESENTATION:**

**Ascites:** 0.5 ml Lyophilized. Reconstitute with 0.5 ml of distilled water.

**Purified:** Purified IgG buffered in PBS and 0.02% NaN<sub>3</sub>. (Purified from ascitic fluid via Protein G Chromatography).

**Biotin/FITC/PE:** Biotin/FITC/PE conjugated IgG buffered in PBS, 0.02% NaN<sub>3</sub> and EIA grade BSA as a stabilizing protein to bring total protein concentration to 4-5 mg/mL.

**STORAGE/STABILITY:**

Lyophilized and reconstituted **ascites** should be stored at -20°C. Reconstituted ascites should be prepared as single use aliquots prior to freezing to minimize freeze/thaw cycles.

For all formats except ascites, store at 4°C. **DO NOT FREEZE PE** conjugates. For long term storage (**Purified, Biotin, FITC, No Azide**), aliquot and freeze unused portion at -20°C in volumes appropriate for single usage. Avoid freeze/thaw cycles.

**SPECIFICATIONS:**

Clone: 34-1-2s

Hybridoma Production:

Immunization: Immunogen: BDF splenocytes  
Donor: C3H spleen

Fusion Partner: Sp2/0-Ag14

Specificity: Mouse H-2K<sup>d</sup>D<sup>d</sup>

*Continued Overleaf.....*

Visit our website for your local distributor.

**CEDARLANE**<sup>®</sup>



[www.cedarlanelabs.com](http://www.cedarlanelabs.com)

An ISO 9001:2000 and ISO 13485:2003  
registered company.

In CANADA: **Toll Free: 1-800-268-5058**

4410 Paletta Court, Burlington, ON L7L 5R2 ph: (289) 288-0001, fax: (289) 288-0020  
e-mail: [general@cedarlanelabs.com](mailto:general@cedarlanelabs.com)

In the USA: **Toll Free: 1-800-721-1644**

1210 Turrentine Street, Burlington, NC 27215 ph: (336) 513-5135, fax: (336) 513-5138  
e-mail: [service@cedarlanelabs.com](mailto:service@cedarlanelabs.com)

Strains Tested: BALB/c, C57BL/6, C3H/He, DBA/1, SJL, B10.M, A.TH, A.TL, B10.A(3R), P/J

Positive: BALB/c, DBA/1, SJL, A.TH, A.TL, B10.A(3R)

Negative: C3H/He, C57BL/6 (+/-), B10.M (+/-), P/J (+/-)

**For a more detailed strain distribution - see reference 1.**

**TEST RESULTS:**


Distribution by Flow Cytometry Analysis:

Mouse Strain: BALB/c

Cell Concentration:  $1 \times 10^6$  cells per test

Antibody Concentration Used:  $0.5 \mu\text{g}/10^6$  cells

<u>Cell Source</u>	<u>Percentage of cells stained above control:</u>
Thymus	82.5%
Spleen	95.7%
Lymph Node	100%



Cell Source: Spleen

Percentage of cells stained above control: 95.7%

**N.B. Appropriate control samples should always be included in any labeling studies.**

**\* For optimal results in various applications, it is recommended that each investigator determine dilutions appropriate for individual use.**

**REFERENCES:**

- 1) Ozato Keiko, Mayer Nanci M., Sachs D.H. Monoclonal antibodies to mouse major histocompatibility complex antigens. Transplantation 34:113-118, 1982.

**FOR RESEARCH USE ONLY**

® is a registered trademark of Cedarlane