



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Agrisera

This product is for research use only (not for diagnostic or therapeutic use)

contact: support@agrisera.com

Agrisera AB | Box 57 | SE-91121 Vännäs | Sweden | +46 (0)935 33 000 | www.agrisera.com

Product no **AS14 2761**

Kelch repeat protein (*Chlamydomonas*)


Product information

Immunogen	Recombinant Cr-Klech, UniProt: A8J2W4 , Locus name: g2987 (from <i>Chlamydomonas Reinhardtii</i>)
Host	Rabbit
Clonality	Polyclonal
Purity	Serum
Format	Lyophilized
Quantity	50 µl
Reconstitution	for reconstitution add 50 µl of sterile water.
Storage	Store lyophilized/reconstituted at -20°C; once reconstituted make aliquots to avoid repeated freeze-thaw cycles. Please, remember to spin tubes briefly prior to opening them to avoid any losses that might occur from lyophilized material adhering to the cap or sides of the tubes.

Application information

Recommended dilution	1:1000 (WB)
Expected apparent MW	33.5 kDa
Confirmed reactivity	<i>Chlamydomonas reinhardtii</i> (strain 21grm cw15, M10), <i>Gonium pectorale</i> , <i>Eudorina elegans</i>
Predicted reactivity	algae
	Species of your interest not listed? Contact us
Not reactive in	no confirmed exceptions from predicted reactivity known in the moment

Application example



Total protein from *Chlamydomonas reinhardtii* (1), *Gonium Pectorale* (2), *Eudorina elegans* (3) was extracted using 1x Laemmli buffer + DTT + protease and phosphatase inhibitors without bromophenol blue heated at 95 oC for ten minutes then vortexed with zirconia beads at max speed for ten minutes. Lysate was spun at 4 oC for ten minutes at max speed in a tabletop centrifuge. Proteins were precipitated out of supernatant in 80% acetone at -20 oC for 20 min, spun at 4 oC at max speed for 10 min. Pellet was washed 1x with 80% acetone and resuspended in 1x LDS sample buffer + DTT + inhibitors at a concentration of 100 mg chlorophyll/µL. Protein equivalent to 1 ug of chlorophyll was loaded on a 4-20% Bis-Tris gel and blotted to nitrocellulose using wet transfer methods. Blot was blocked with 5% milk in TBS-T for 1hr/RT. Blot was incubated in the primary antibody at a dilution of 1:1000 in 5% milk in TBS-T ON/4oC. The antibody solution was decanted and the blot was rinsed 3 times for 5 min in TBS-T at RT with agitation. Blot was incubated in Agrisera matching secondary diluted to 1:25,000 in 5% milk in TBS-T for 1hr/RT with agitation. The blot was washed as above and developed with chemiluminescent detection reagent and imaged on a Licor C-Digit blot scanner.

Agrisera

This product is **for research use only** (not for diagnostic or therapeutic use)

contact: support@agrisera.com

Agrisera AB | Box 57 | SE-91121 Vännäs | Sweden | +46 (0)935 33 000 | www.agrisera.com

Exposure time was 12 min.

Courtesy of Jessica Rakijas, Kansas State University, USA