



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Technically  
Speaking

**CEDARLANE**<sup>®</sup>   
[www.cedarlanelabs.com](http://www.cedarlanelabs.com)

Conveniently Delivering You Today's Innovations  
for the Science of Tomorrow™

## Glutaraldehyde Stabilized Turkey RBCs (10%)

**CLD118**  
Lot: 1624

### DESCRIPTION:

Cedarlane's **CLD118** are Glutaraldehyde-stabilized freshly prepared Turkey Red Blood Cells. Glutaraldehyde stabilized red blood cells are produced from freshly washed turkey RBCs by brief exposure to a glutaraldehyde-buffer solution and exhaustive washing in saline. This treatment largely preserves the native antigenicity of the erythrocytes while rendering the cells generally resistant to lysis by osmotic shock, freeze-thawing or immune hemolysis. This reagent can be used directly in hemagglutination procedures, or may be coupled with various proteins for hemagglutination testing.

### PRESENTATION:

20 ml of glutaraldehyde-stabilized turkey RBCs in saline with 0.1% NaN<sub>3</sub> (10%).

### STORAGE:

Store at 4°C. DO NOT FREEZE.

### APPLICATIONS:

This product is suitable for use in hemagglutination assays.

### TEST TYPE:

#### Settling Pattern:

100 µl of 2% resuspended cells of lot 1624 when mixed with 100 µl of saline in U-bottom microtiter plate wells settle into buttons within 2-4 hours.

#### Presence of Agglutinogens:

100 µl of 2% resuspended cells of lot 1624 when mixed with 100 µl of agglutinin dilutions made in 1:100 diluted normal rabbit serum in saline in U-bottom microtiter plate wells; incubated @ 37°C for 20 minutes and 20°C for 2-4 hours form typical hemagglutination patterns.

#### Microscope Observations:

No crenation; Single, Entire Cells

#### Osmotic Shock Test:

No lysis in 100% distilled water

**Optimum conditions for use of this product in analytical procedures should be established in each laboratory**

**Laboratory Reagent For Research Use Only**

XT 01/14/16

Visit our website for your local distributor.

**CEDARLANE**<sup>®</sup> 

[www.cedarlanelabs.com](http://www.cedarlanelabs.com)

An ISO 9001:2000 and ISO 13485:2003  
registered company.

In CANADA: Toll Free: 1-800-268-5058

4410 Paletta Court, Burlington, ON L7L 5R2 ph: (289) 288-0001, fax: (289) 288-0020  
e-mail: [general@cedarlanelabs.com](mailto:general@cedarlanelabs.com)

In the USA: Toll Free: 1-800-721-1644

1210 Turrentine Street, Burlington, NC 27215 ph: (336) 513-5135, fax: (336) 513-5138  
e-mail: [service@cedarlanelabs.com](mailto:service@cedarlanelabs.com)