



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Technically
Speaking

CEDARLANE[®]
www.cedarlanelabs.com



Conveniently Delivering You Today's Innovations
for the Science of Tomorrow™

Recombinant Human Angiotensin Converting Enzyme 2 (19-740 a.a.)

CLENZ1122
CLENZ1122-2
CLENZ1122-3

Introduction: ACE2 (Angiotensin converting enzyme 2) an enzyme bound to cell membranes in various organs such as intestines, arteries, lungs, heart & kidney. ACE2 an entry receptor of SARS coronaviruses as well as SARS-CoV-2. The coronavirus spike (S) glycoprotein is a class I viral fusion antigen located on the external envelope of the virion that takes part in a critical part in viral infection by identifying host cell receptors and facilitating fusion of the viral and cellular membranes. 2 main domains in coronavirus S1 have been recognized, the N-terminal domain and C-terminal domain. One or the other and/or both S1 domains function as a receptor-binding domain. SARS-CoV + MERS-CoV equally use C-domain to attach their receptors. ACE2 is a type I transmembrane antigen with an extracellular N-terminal domain having the catalytic site and an intracellular C-terminal tail. ACE2 obtains a signal peptide, a transmembrane domain, and a single metalloproteinase active site containing an HEXXH zinc-binding domain. ACE-2 plays a role as a mono-carboxypeptidase which degrades Ang I to produce the nonapeptide Ang 1-9 and Ang II to create the heptapeptide Ang 1-7.

Description: The CHO derived ACE2 Human recombinant protein contains the extracellular domain amino acids 19-740 fused to Fc tag at C-terminal. ACE2 Protein binds to SARS Coronavirus-2 [CoV-2019] Spike receptor binding domain.

Source: CHO Cells.

Purification Method: Purified by Protein-G chromatographic technique.

Presentation: 1 µg (CLENZ1122), 5 µg (CLENZ1122-2), or 10 µg (CLENZ1122-3), sterile filtered clear solution. ACE2 Human protein solution is supplied in 50mM Tris-HCl, pH7.5, and 90mM glycine.

Stability: Store at -20°C.

Purity: Greater than 95.0% as determined by SDS-PAGE.

Biological Activity: ACE2 activity was measured by its binding ability in a functional ELISA. The immobilized Recombinant Human ACE2 protein binds to SARS CoV2 Spike protein Receptor Binding Domain at 2 µg per ml.

Laboratory Reagent For Research Use Only

MW 04/20/20

Visit our website for your local distributor.

CEDARLANE[®]



www.cedarlanelabs.com

An ISO 9001:2015 and ISO 13485:2016
registered company.

In CANADA: Toll Free: 1-800-268-5058

4410 Paletta Court, Burlington, ON L7L 5R2 ph: (289) 288-0001, fax: (289) 288-0020
e-mail: general@cedarlanelabs.com

In the USA: Toll Free: 1-800-721-1644

1210 Turrentine Street, Burlington, NC 27215 ph: (336) 513-5135, fax: (336) 513-5138
e-mail: service@cedarlanelabs.com