



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

TECHNICALLY *Speaking*

Place your order with CEDARLANE® or your local distributor.

Please contact CEDARLANE® for lot specific information.

Anti-Human HLA-DR Ascites Monoclonal Antibody

CLHLA-03A

LOT: 0311

DESCRIPTION:

Cedarlane's anti-human HLA-DR monoclonal antibody recognizes the HLA-DR (MHC class II) antigen.

Applications include: flow cytometry, cryostat sections, Parafin-embedded sections

PRESENTATION:

0.5 mL, lyophilized ascites.

STORAGE/STABILITY:

Lyophilized form stable at -20°C for long term storage or 4°C for short term storage. Reconstitute with 0.5 ml of cold distilled water. For long term storage, aliquot and freeze unused portions at -20°C in volumes appropriate for single usage. Avoid repeated freeze/thaw cycles as this may denature the antibody.

For more information or to place an order please contact...

CEDARLANE®
LABORATORIES LIMITED



*toll free: 1-800-268-5058
in North America*

phone: (905) 878-8891 • fax: (905) 878-7800

5516 - 8th Line, R.R.#2, Hornby, Ontario, CANADA L0P 1E0

or visit our website for a list of our international distributors including contact information
website: www.cedarlanelabs.com • e-mail: info@cedarlanelabs.com

SPECIFICATIONS:

Clone: YD1/63.4.10

Hybridoma Production:

Immunization: Immunogen: DAUDI cells
Donor: immunized DA rat spleen cells

Fusion Partner: Y3 Ag1.2.3 rat myeloma

Specificity: HLA-DR antigen

Ig Class: Rat IgG_{2a}

Format: Ascitic fluid filtered to 0.45 µm (non-sterile) and lyophilized

FLOW CYTOMETRY ANALYSIS:

Method:

1. Prepare a cell suspension in media A. For cell preparations, deplete the red blood cell population with Lympholyte[®]-H (CL5010) cell separation medium.
2. Wash 2 times.
3. Resuspend the cells to a concentration of 2×10^7 cells/ml in media A. Add 50 µl of this suspension to each tube (each tube will then contain 1×10^6 cells, representing 1 test).
4. To each tube, add 50 µl of a 1/10000-1/20000 dilution of CLHLA-03A per 10^6 cells.
5. Vortex the tubes to ensure thorough mixing of antibody and cells.
6. Incubate the tubes for 30 minutes at 4°C.
7. Wash 2 times at 4°C.
8. Add 100 µl of secondary antibody **CLCC40001** (FITC Goat anti-rat IgG) at a 1:500 dilution.
9. Incubate the tubes at 4°C for 30-60 minutes.
(It is recommended that the tubes are protected from light since most fluorochromes are light sensitive).
10. Wash 2 times at 4°C in media B.
11. Resuspend the cell pellet in 50 µl ice cold media B.
12. Transfer to suitable tubes for flow cytometric analysis containing 15 µl of propidium iodide at 0.5 mg/ml in PBS. This stains dead cells by intercalating in DNA.

Media:

- A. Phosphate buffered saline (pH 7.2) + 5% normal serum of host species + sodium azide (100 µl of 2M sodium azide in 100 mls).
- B. Phosphate buffered saline (pH 7.2) + 0.5% Bovine serum albumin + sodium azide (100 µl of 2M sodium azide in 100 mls).


Results:

Tissue Distribution by Flow Cytometry Analysis:

Cell Concentration: 1×10^6 cells per test

Antibody Concentration Used: 1/20000

Isotypic Control: Purified Rat IgG_{2a} (CLCR2a00)



Cell Source: Peripheral Blood Lymphocytes
Percentage of cells stained above control: 10.6%

N.B. Appropriate control samples should always be included in any labeling studies.

*** For optimal results in various applications, it is recommended that each investigator determine dilutions appropriate for individual use.**

REFERENCES:

1. Janossy, G., Tidman, N., Crawford, D., Papageorgiou, E. S., Prentice, H. G., Francis, G., Bradstock, K. F., McConnell, I., Secher, D., and Milstein, C. (1980). Protides of Biol. Fluids 28: 523-528

FOR RESEARCH USE ONLY

® is a Registered Trademark of Cedarlane Laboratories Limited.

JB/18/06/01