



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 



Place your order with CEDARLANE® or your local distributor.

*Please contact CEDARLANE® for lot specific information.*

**Purified Anti-Human HLA-DR  
Monoclonal Antibody**

**CLHLA-03AP**

**LOT: 0321**

**DESCRIPTION:**

Cedarlane's purified anti-human HLA-DR monoclonal antibody recognizes the HLA-DR (MHC class II) antigen.

Applications include: flow cytometry, cryostat sections, Parafin-embedded sections

**PRESENTATION:**

500µg purified Ig buffered in PBS with 0.02% NaN<sub>3</sub>.

**STORAGE/STABILITY:**

Store at 4°C. For long term storage, aliquot and freeze unused portion at -20°C in volumes appropriate for single usage. Avoid repeated freeze/thaw cycles as this may denature the antibody.

For more information or to place an order please contact...

**CEDARLANE®**  
**LABORATORIES LIMITED**



*toll free: 1-800-268-5058  
in North America*

phone: (905) 878-8891 • fax: (905) 878-7800

5516 - 8th Line, R.R.#2, Hornby, Ontario, CANADA L0P 1E0

or visit our website for a list of our international distributors including contact information  
website: [www.cedarlanelabs.com](http://www.cedarlanelabs.com) • e-mail: [info@cedarlanelabs.com](mailto:info@cedarlanelabs.com)

**SPECIFICATIONS:**

Clone: YD1/63.4.10

Hybridoma Production:

Immunization: Immunogen: DAUDI cells

Donor: immunized DA rat spleen cells

Fusion Partner: Y3 Ag1.2.3 rat myeloma

Specificity: HLA-DR antigen

Ig Class: Rat IgG<sub>2a</sub>

Format: Purified Ig buffered in PBS and 0.02% NaN<sub>3</sub> (purified from ascites via Protien G Chromatography).

Antibody Concentration: 1.0 mg/ml

**FLOW CYTOMETRY ANALYSIS:**

Method:

1. Prepare a cell suspension in media A. For cell preparations, deplete the red blood cell population with Lympholyte<sup>®</sup>-H (CL5010) cell separation medium.
2. Wash 2 times.
3. Resuspend the cells to a concentration of  $2 \times 10^7$  cells/ml in media A. Add 50  $\mu$ l of this suspension to each tube (each tube will then contain  $1 \times 10^6$  cells, representing 1 test).
4. To each tube, add 50  $\mu$ l of a 0.2  $\mu$ g – 0.05  $\mu$ g dilution of CLHLA-03AP per  $10^6$  cells.
5. Vortex the tubes to ensure thorough mixing of antibody and cells.
6. Incubate the tubes for 30 minutes at 4°C.
7. Wash 2 times at 4°C.
8. Add 100  $\mu$ l of secondary antibody **CLCC40001** (FITC Goat anti-rat IgG) at a 1:500 dilution.
9. Incubate the tubes at 4°C for 30-60 minutes.  
(It is recommended that the tubes are protected from light since most fluorochromes are light sensitive).
10. Wash 2 times at 4°C in media B.
11. Resuspend the cell pellet in 50  $\mu$ l ice cold media B.
12. Transfer to suitable tubes for flow cytometric analysis containing 15  $\mu$ l of propidium iodide at 0.5 mg/ml in PBS. This stains dead cells by intercalating in DNA.

Media:

- A. Phosphate buffered saline (pH 7.2) + 5% normal serum of host species + sodium azide (100  $\mu$ l of 2M sodium azide in 100 mls).
- B. Phosphate buffered saline (pH 7.2) + 0.5% Bovine serum albumin + sodium azide (100  $\mu$ l of 2M sodium azide in 100 mls).


Results:Tissue Distribution by Flow Cytometry Analysis:

Cell Concentration:  $1 \times 10^6$  cells per test

Antibody Concentration Used: 0.05  $\mu$ g

Isotypic Control: Purified Rat IgG<sub>2a</sub> (CLCR2a00)

Cell Source: Peripheral Blood Lymphocytes



Percentage of cells stained above control: 9.77%

**N.B. Appropriate control samples should always be included in any labeling studies.**

**\* For optimal results in various applications, it is recommended that each investigator determine dilutions appropriate for individual use.**

**REFERENCES:**

1. Janossy, G., Tidman, N., Crawford, D., Papageorgiou, E. S., Prentice, H. G., Francis, G., Bradstock, K. F., McConnell, I., Secher, D., and Milstein, C. (1980).  
Protides of Biol. Fluids 28: 523-528

**FOR RESEARCH USE ONLY**

® is a Registered Trademark of Cedarlane Laboratories Limited.

JB/18/06/01