



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Technically  
Speaking

**CEDARLANE**<sup>®</sup>   
[www.cedarlanelabs.com](http://www.cedarlanelabs.com)

Conveniently Delivering You Today's Innovations  
for the Science of Tomorrow™

## Bovine Endothelial Mitogen

**CLPRO603**  
**CLPRO603-2**  
**CLPRO603-3**  
**Lot:**

**Description:** Endothelial Mitogen can be used for Human & Mammalian Vascular Endothelial Cells at concentration of 0.1-0.3mg/ml of serum supplemented media & 0.01-0.1mg with heparin. Endothelial Mitogen can be used for Hybridoma Cell Cloning at concentrations of 0.025-0.15mg/ml of serum supplemented media. Endothelial Mitogen can be used for Balb/C-3T3 Cells at concentrations of 0.2mg/ml of serum Supplemented media.

**Source:** Bovine Hypothalamus.

**Presentation:** 5 mg (**CLPRO603**), 20 mg (**CLPRO603-2**), or 100 mg (**CLPRO603-3**), sterile filtered Endothelial-Mitogen was lyophilized from a concentrated solution containing 0.1M NaCl and 5mM NaH<sub>2</sub>PO<sub>4</sub>, pH 7.4.

**Solubility:** It is recommended to reconstitute the lyophilized Endothelial-Mitogen Bovine Recombinant in sterile 18MΩ-cm H<sub>2</sub>O not less than 100µg/ml, which can then be further diluted to other aqueous solutions.

**Stability:** Lyophilized Endothelial-Mitogen Bovine although stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon reconstitution Endothelial-Mitogen Bovine Recombinant should be stored at 4°C between 2-7 days and for future use below -18°C. Please prevent freeze-thaw cycles.

**Laboratory Reagent For Research Use Only**

JK 01/29/13

Visit our website for your local distributor.

**CEDARLANE**<sup>®</sup> 

[www.cedarlanelabs.com](http://www.cedarlanelabs.com)

An ISO 9001:2000 and ISO 13485:2003  
registered company.

In CANADA: **Toll Free: 1-800-268-5058**

4410 Paletta Court, Burlington, ON L7L 5R2 ph: (289) 288-0001, fax: (289) 288-0020  
e-mail: [general@cedarlanelabs.com](mailto:general@cedarlanelabs.com)

In the USA: **Toll Free: 1-800-721-1644**

1210 Turrentine Street, Burlington, NC 27215 ph: (336) 513-5135, fax: (336) 513-5138  
e-mail: [service@cedarlanelabs.com](mailto:service@cedarlanelabs.com)