



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Technically
Speaking

CEDARLANE[®] 
www.cedarlanelabs.com

Conveniently Delivering You Today's Innovations
for the Science of Tomorrow™

Mouse Anti-Human B-Cell, CD20 Monoclonal Antibody

CLSG36516
CLSG36516-05
Lot:

Intended Use: This product is intended for qualitative immunohistochemistry with normal and neoplastic formalin-fixed, paraffin-embedded tissue sections, to be viewed by light microscopy. Clinical interpretation of staining results should be accompanied by histological studies with proper controls. Patients' clinical histories and other relevant diagnostic tests should be utilized by a qualified person(s) when evaluating and interpreting results.

Immunogen: BALB/C mice were injected with human tonsil B cells.

Clone: L26

Isotype: IgG2a, Kappa

Format: 1 ml (CLSG36516) or 0.5 ml (CLSG36516-05). This antibody is supplied as tissue culture supernatant containing sodium azide as a preservative.

Titer/Working Dilution: This antibody may be diluted to a titer of 1:25-1:50 using the PolyVue Detection System. The final dilution should be determined by the user based upon the staining conditions employed.

Staining Protocol: We suggest an incubation period of thirty minutes at room temperature. Optimal incubation conditions should be determined by the user based upon the fixation conditions and staining system employed. Suitable for formalin fixed, paraffin embedded tissue sections.

Specificity: This antibody is specific to a 33 kD polypeptide present on the majority of B cells in peripheral blood and lymphoid tissue. No reactivity with other hematopoietic cells has been observed.

Positive Control: Tonsil

Cellular Localization: Cell membrane and cytoplasm

Continued...

Visit our website for your local distributor.

CEDARLANE[®] 

www.cedarlanelabs.com

An ISO 9001:2000 and ISO 13485:2003
registered company.

In CANADA: Toll Free: 1-800-268-5058

4410 Paletta Court, Burlington, ON L7L 5R2 ph: (289) 288-0001, fax: (289) 288-0020
e-mail: general@cedarlanelabs.com

In the USA: Toll Free: 1-800-721-1644

1210 Turrentine Street, Burlington, NC 27215 ph: (336) 513-5135, fax: (336) 513-5138
e-mail: service@cedarlanelabs.com

Storage:

Store the antibody at 2-8°C. Do not use it beyond the expiration date stated on the label.

References:

- i) Ishii et al. Clin Exp Immunol 58: 183, 1984.
- ii) Mason et al. Am J Pathol 136: 1215, 1990.
- iii) Cartun et al. Am J Pathol 129: 415,1987.

BA 10/25/12

Laboratory Reagent For Research Use Only.