



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

# TECHNICALLY *Speaking*

Place your order with CEDARLANE® or your local distributor.

Please contact CEDARLANE® for lot specific information.

## Purified Mouse anti-Human Caspase 3 Monoclonal Antibody

CLSG36537  
CLSG36537-05

Lot:

**Immunogen:** BALB/C mice were injected with recombinant full length human Caspase 3 protein.

**Clone:** 3CSP03

**Isotype:** IgG2a

**Format:** 1 ml (CLSG37537) or 0.5 ml (CLSG36537-05) of purified immunoglobulin fraction containing sodium azide as preservative.

**Titer/Working Dilution:** This antibody may be diluted to a titer of 1:25-1:75 in an ABC method. However, depending on the staining conditions employed, we suggest that final dilution should be determined by the user.

**Staining Protocol:** We suggest an incubation period of 30 minutes at room temperature. However, depending upon the fixation conditions and the staining system employed, optimal incubation should be determined by the user. Formalin fixed paraffin embedded tissue sections require high temperature antigen unmasking with 10 mM citrate buffer, pH 6.0 prior to immunostaining.

**Specificity:** This antibody is specific to a 32 kDa protein, which is identified as Caspase 3. Caspase 3 is associated with induction of apoptosis. Caspase 3 is ubiquitously express and is synthesized as an inactive proenzyme.

Continued Overleaf...

For more information or to place an order please contact...

**CEDARLANE®**  
**LABORATORIES LIMITED**



**toll free: 1-800-268-5058**  
**in North America**

phone: (905) 878-8891 • fax: (905) 878-7800

5516 - 8th Line, R.R.#2, Hornby, Ontario, CANADA L0P 1E0

or visit our website for a list of our international distributors including contact information  
website: [www.cedarlanelabs.com](http://www.cedarlanelabs.com) • e-mail: [info@cedarlanelabs.com](mailto:info@cedarlanelabs.com)

**Positive Control:** Tonsil

**Cellular Localization:** Cytoplasmic

**Other Applications:** Immunoprecipitation, Western Blotting

**Storage:** Store the antibody at 4-8°C. Do not use it beyond the expiration date stated on the label.

**References:**

- i) Krajewska et al. Cancer Res 57 (8): 1605, 1997.
- ii) Mallat et al. Circulation 96 (2): 424, 1997

**Laboratory Reagent For Research Use Only**

JV05/14/07