



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

TECHNICALLY Speaking

Place your order with CEDARLANE® or your local distributor.

Please contact CEDARLANE® for lot specific information.

Mouse Anti-Human CD10 antigen, CALLA Monoclonal Antibody

CLSG36560

CLSG36560-05

Lot:

Immunogen:	Prokaryotic recombinant protein corresponding to the external domain of the CD10 glycoprotein.
Clone:	56C6
Isotype:	IgG1
Format:	1 ml (CLSG36560) or 0.5 ml (CLSG36560-05). This antibody is supplied as tissue culture supernatant containing sodium azide as preservative.
Titer/Working Dilution:	This antibody may be diluted to a titer of 1:20-1:30 in an ABC method. However, depending on the staining conditions employed, we suggest that final dilution should be determined by the user.
Staining Protocol:	We suggest an incubation period of 30 minutes at room temperature. However, depending upon the fixation conditions and the staining system employed, optimal incubation should be determined by the user. <u>High temperature antigen unmasking is required prior to immunostaining of formalin fixed paraffin embedded tissue sections.</u>
Specificity:	This antibody is specific to human CD10 antigen of 100 kDa, also known as common acute lymphocytic leukemia antigen (CALLA). CD10 antigen has been identified on the surface of early lymphoid progenitor cells, immature B cells with in adult bone marrow and germinal center B cells with in lymphoid tissue.
Positive Control:	Tonsil
Cellular Localization:	Cell membrane
Storage:	Store the antibody at 2-8°C. Do not use beyond expiration date stated on the label.
References:	i) Carrel et al. Melanoma Res 3: 319, 1993. ii) Haralambidou et al. J Clin Pathol 40: 490, 1987. iii) Kiyokawa et al. Clin Exp Immunol 79: 322, 1990. iv) Scheuermann et al. Leukemia and Lymphoma 18: 385, 1995.

Laboratory Reagent For Research Use Only

JV 02/19/08

For more information or to place an order please contact...

CEDARLANE®
LABORATORIES LIMITED



toll free: 1-800-268-5058

in North America

phone: (905) 878-8891 • fax: (905) 878-7800

5516 - 8th Line, R.R.#2, Hornby, Ontario, CANADA L0P 1E0

or visit our website for a list of our international distributors including contact information

website: www.cedarlanelabs.com • e-mail: info@cedarlanelabs.com